arXiv:0908.3580v2 [math.AT] 24 Oct 2009

ON HOMOTOPY GROUPS OF THE SUSPENDED CLASSIFYING SPACES

ROMAN MIKHAILOV AND JIE WU

Abstract. In this paper, we determine the homotopy groups $\pi_4(\Sigma K(A,1))$ and $\pi_5(\Sigma K(A,1))$ for abelian groups A by using different facts and methods from group theory and homotopy theory: derived functors, the Carlsson simplicial construction, the Baues-Goerss spectral sequence, homotopy decompositions and the methods of algebraic K-theory. As the applications, we also determine $\pi_i(\Sigma K(G,1))$ with $i=4,5$ for some non-abelian groups $G=\Sigma_3$ and SL($\mathbb{Z}$), and $\pi_4(\Sigma K(A_4,1))$ for the 4-th alternating group A_4.

1. Introduction

It is well-known that the suspension functor applied to a topological space shifts homology groups, but "chaotically" changes homotopy groups. For example, one can take a circle S^1, whose homotopy type is very simple. Its suspension $\Sigma S^1 = S^2$ has obvious homology groups, however the problem of investigating the homotopy groups of S^2 is one of the deepest problems of algebraic topology. Consider the following functors from the category of groups to the category of abelian groups:

$$\pi_n(\Sigma^m K(-,1)) : \mathsf{Gr} \to \mathsf{Ab},\ n \geq 1, m \geq 1$$

defined by $A \mapsto \pi_n(\Sigma^m K(A,1))$, where Σ^m is the m-fold suspension. It is clear that $\pi_n(\Sigma^m K(\mathbb{Z},1)) = \pi_n(S^{m+1})$, that is the homotopy groups of spheres appear as the simplest case of a general theory of homotopy groups of suspensions of classifying spaces.

For the case $m=1,2$ and $n=3,4$ there is the following natural commutative diagram with exact rows [6]:

$$\begin{tikzcd}
0 \arrow[r] & \pi_3(\Sigma K(G,1)) \arrow[r] \arrow[d, two heads] & G \otimes G \arrow[r] \arrow[d, two heads] & {[G,G]} \arrow[r] \arrow[d, equal] & 1 \\
0 \arrow[r] & \pi_4(\Sigma^2 K(G,1)) \arrow[r] \arrow[d, two heads] & G \widetilde{\otimes} G \arrow[r] \arrow[d, two heads] & {[G,G]} \arrow[r] \arrow[d, equal] & 1 \\
0 \arrow[r] & H_2(G) \arrow[r] & G \wedge G \arrow[r] & {[G,G]} \arrow[r] & 1
\end{tikzcd} \tag{1.1}$$

1991 *Mathematics Subject Classification.* Primary 55Q52; Secondary 55P20, 55P40, 55P65, 55Q35.

Key words and phrases. homotopy groups, Whitehead exact sequence, spectral sequences, Moore spaces, suspensions of $K(G,1)$ spaces, simplicial group.

The research of the first author is partially supported by RFBR (grant 08-01-91300 IND_a). The research of the second author is partially supported by the Academic Research Fund of the National University of Singapore R-146-000-101-112.

where $G\otimes G$ is the non-abelian square of G in the sense of Brown-Loday [6], $G\tilde{\otimes}G$ (resp. $G\wedge G$) is the quotient of $G\otimes G$ by the normal subgroup generated by elements $g\otimes h+h\otimes g$ (resp. $g\otimes g, g\in G$). In particular, for an abelian group A, there are natural isomorphisms

$$\begin{aligned}\pi_3(\Sigma K(A,1)) &\simeq A\otimes A\\ \pi_4(\Sigma^2 K(A,1)) &\simeq \pi_2^S K(A,1)\simeq A\tilde{\otimes}A.\end{aligned}$$

The purpose of this article is to determine the homotopy groups $\pi_4(\Sigma K(A,1))$ and $\pi_5(\Sigma K(A,1))$ for abelian groups A. In order to investigate the structure of these homotopy groups, we use different facts and methods of group theory and homotopy theory: derived functors, the Carlsson simplicial construction, the Baues-Goerss spectral sequence [4], homotopy decompositions and the methods of algebraic K-theory. The combination of these different methods provides an effective way for determining these homotopy groups. As reader will see, some our computations use commutator tricks in simplicial groups.

The homotopy group $\pi_4(\Sigma K(A,1))$ as a functor on A can be given as follows:

Theorem 1.1 (Theorem 3.1). *Let A be any abelian group. Then there is a natural short exact sequence*

$$(\Lambda^2(A)\otimes A)^{\oplus 2}\oplus A\otimes A\otimes \mathbb{Z}/2 \hookrightarrow \pi_4(\Sigma K(A,1)) \twoheadrightarrow \mathrm{Tor}(A,A).$$

Moreover $(\Lambda^2(A)\otimes A)^{\oplus 2}$ is an (unnatural) summand of $\pi_4(\Sigma K(A,1))$.

An interesting point of this theorem is that the functor $\pi_4(\Sigma K(A,1))$ has $\mathrm{Tor}(A,A)$ as a natural quotient. For determining the structure of the group $\pi_4(\Sigma K(A,1))$, one has to solve the group extension problem in Theorem 1.1. For finitely generated abelian groups A, we are able to solve this problem. Given a finitely generated abelian group A, let

$$A = A_1\oplus \bigoplus_{\substack{r\geq 1\\ p \text{ is a prime}}} A_{p^r}$$

be the primary decomposition of A, where A_1 is torsion free and A_{p^r} is a free $\mathbb{Z}/p^r$-module.

Theorem 1.2 (Theorem 3.2). *Let A be any finitely generated abelian group. Let $A = A_2\oplus B$ with $B=A_1\oplus\bigoplus_{p^r\neq 2}A_{p^r}$. Then*

$$\begin{aligned}\pi_4(\Sigma K(A,1)) \;\cong\; &\tfrac{1}{2}(A_2\otimes A_2)\oplus(A_2\otimes B)^{\oplus 2}\oplus B^{\otimes 2}\otimes\mathbb{Z}/2\oplus(A\otimes\Lambda^2(A))^{\oplus 2}\\ &\oplus\mathrm{Tor}(A_2,B)^{\oplus 2}\oplus\mathrm{Tor}(B,B),\end{aligned}$$

where $\frac{1}{2}(A_2\otimes A_2)$ is a free $\mathbb{Z}/4$-module with rank of $\dim_{\mathbb{Z}/2}(A_2\otimes A_2)$.

One point of this theorem is that the (maximal) elementary 2-group summand A_2 of A plays a key role in the group extension problem. Roughly speaking $A_2\otimes A_2$ is half down in the group $\pi_4(\Sigma K(A,1))$.

As the applications of Theorems 1.1 and 1.2, we are able to compute $\pi_4(M(\mathbb{Z}/2^r,2))$ and their connections with $\pi_4(\Sigma K(\mathbb{Z}/2^r,1))$. As the direct consequences, the homotopy groups $\pi_4(\Sigma\mathbb{R}\mathrm{P}^n)$ and $\pi_4(\Sigma K(\Sigma_3,1))$ are determined. (See subsection 3.2 for the computations of these homotopy groups.)

For the homotopy group $\pi_5(\Sigma K(A,1))$, as a functor, it can be described by two exact sequences given in diagram (4.1). Unfortunately it seems too complicated to produce a canonical functorial short exact sequence description for the functor $\pi_5(\Sigma K(A,1))$ from diagram (4.1). For any finitely generated abelian group A, we determine $\pi_5(\Sigma K(A,1))$ in an un-functorial way by the following steps:

1) From the Hopf fibration, $\pi_5(\Sigma K(A,1)) \cong \pi_5(\Sigma K(A,1) \wedge K(A,1))$;
2) Take a primary decomposition of A and write $K(A,1)$ as a product of copies of $S^1 = K(\mathbb{Z},1)$ and $K(\mathbb{Z}/p^r,1)$;
3) By using the fact that

$$\Sigma X \times Y \simeq \Sigma X \vee \Sigma Y \vee \Sigma X \wedge Y,$$

write $\Sigma K(A,1) \wedge K(A,1)$ as a wedge of the spaces in the form

$$X = \Sigma^m K(\mathbb{Z}/p_1^{r_1},1) \wedge K(\mathbb{Z}/p_2^{r_2},1) \wedge \cdots \wedge K(\mathbb{Z}/p_t^{r_t},1)$$

with $m + t \geq 3$ and $m \geq 1$;
4) By applying the Hilton-Milnor Theorem, $\pi_5(\Sigma K(A,1))$ becomes a summation of $\pi_5(X)$ for some X in the above form.

For the spaces X in the above form, it is contractible if $p_i \neq p_j$ for some $i \neq j$ and $\pi_5(X)$ can be determined in Proposition 4.2 for an odd prime p. The only difficult part is to compute $\pi_5(X)$ for X given in the form

$$X = \Sigma^m K(\mathbb{Z}/2^{r_1},1) \wedge \cdots \wedge K(\mathbb{Z}/2^{r_t},1)$$

with $m + t \geq 3$ and $m \geq 1$. Our computations are then given case-by-case (Propositions 4.4-4.6 and Theorems 4.1-4.4), in which different methods are involved. An instructional example is as follows:

Let $A = \mathbb{Z} \oplus \mathbb{Z}/2$. According to 1), $\pi_5(\Sigma K(A,1)) \cong \pi_5(\Sigma K(A,1) \wedge K(A,1))$. As in 3),

$$\begin{array}{rcl} \Sigma K(A,1) \wedge K(A,1) & \simeq & \Sigma(S^1 \times \mathbb{R}\mathrm{P}^\infty) \wedge (S^1 \times \mathbb{R}\mathrm{P}^\infty) \\ & \simeq & \Sigma(S^1 \vee \mathbb{R}\mathrm{P}^\infty \vee \Sigma\mathbb{R}\mathrm{P}^\infty) \wedge (S^1 \vee \mathbb{R}\mathrm{P}^\infty \vee \Sigma\mathbb{R}\mathrm{P}^\infty) \\ & = & S^3 \vee \bigvee^2 \Sigma^2\mathbb{R}\mathrm{P}^\infty \vee \bigvee^2 \Sigma^3\mathbb{R}\mathrm{P}^\infty \vee \Sigma\mathbb{R}\mathrm{P}^\infty \wedge \mathbb{R}\mathrm{P}^\infty \vee \\ & & \vee \bigvee^2 \Sigma^2\mathbb{R}\mathrm{P}^\infty \wedge \mathbb{R}\mathrm{P}^\infty \vee \Sigma^3\mathbb{R}\mathrm{P}^\infty \wedge \mathbb{R}\mathrm{P}^\infty. \end{array}$$

By applying the Hilton-Milnor Theorem as in 4), $\pi_5(\Sigma K(A,1) \wedge K(A,1))$ is a summation of

$$\pi_5(S^3),\ \pi_5(\Sigma^2\mathbb{R}\mathrm{P}^\infty), \pi_5(\Sigma\mathbb{R}\mathrm{P}^\infty \wedge \mathbb{R}\mathrm{P}^\infty), \pi_5(\Sigma(\mathbb{R}\mathrm{P}^\infty)^{\wedge 3}), \cdots$$

with multiplicities. From Theorem 4.4, we have

$$\pi_5(\Sigma^2\mathbb{R}\mathrm{P}^\infty) = \mathbb{Z}/8$$

and by Proposition 4.6 and Theorem 4.1, we have

$$\pi_5(\Sigma\mathbb{R}\mathrm{P}^\infty \wedge \mathbb{R}\mathrm{P}^\infty) = \pi_5(\Sigma(\mathbb{R}\mathrm{P}^\infty)^{\wedge 3}) = \mathbb{Z}/2^{\oplus 2}.$$

The group $\pi_5(\Sigma K(\mathbb{Z} \oplus \mathbb{Z}/2))$ will be determined by filling all possible summands with multiplicities.

As the applications of our computations on $\pi_5(\Sigma K(A,1))$, we are able to determine $\pi_5(\Sigma\mathbb{R}\mathrm{P}^n)$ (Proposition 4.8) and $\pi_5(\Sigma K(\Sigma_3,1))$ (Proposition 4.9).

In section 2 we recall certain facts from the homotopy theory, such as the Whitehead exact sequence, the Carlsson simplicial construction and describe a spectral sequence (2.9), which converges to $\pi_*(\Sigma^m K(A,1))$ for any abelian group A, with E^2-terms are given by the derived functors of certain polynomial functors. We illustrate how it works in Theorem 4.4 for computing

$$\pi_5(\Sigma^2 K(\mathbb{Z}/2^r,1)) = \begin{array}{lll} \mathbb{Z}/8 & \text{if} & r=1, \\ \mathbb{Z}/2^{r+1}\oplus\mathbb{Z}/2 & \text{if} & r>1. \end{array}$$

The interesting point is of course how $\mathbb{Z}/8$ shows up in the case $r=1$ while it becomes $\mathbb{Z}/2^{r+1}\oplus\mathbb{Z}/2$ for $r>1$. The proof is also based on the computations of the derived functors of the antisymmetric square $\tilde{\otimes}^2$.

There is a natural relation between the problem considered and algebraic K-theory. Since the plus-construction $K(G,1)\to K(G,1)^+$ is a homological equivalence, there is a natural weak homotopy equivalence

$$\Sigma K(G,1)\to\Sigma(K(G,1)^+)$$

This defines the natural suspension map:

$$\pi_n(K(G,1)^+)\to\pi_{n+1}(\Sigma(K(G,1)^+))=\pi_{n+1}(\Sigma K(G,1))$$

for $n\geq 1$. This map was studied in [3] in the case of a perfect group G. We consider the case $G=E(R)$, i.e. the group of elementary matrices over a ring R. In this case the natural map

$$K_3(R)=\pi_3(K(E(R),1)^+)\to\pi_4(\Sigma K(E(R),1))$$

is an isomorphism (Theorem 5.1). The natural relation to K-theory gives a way how to compute homotopy groups $\pi_i(\Sigma K(E(R),1))$ for $i=4,5$ for some rings. For example, the case $G=SL(\mathbb{Z})$ is considered. As an application of our methods, we also determine that $\pi_4(\Sigma K(A_4,1))=\mathbb{Z}/4$ for the 4-th alternating group A_4.

The article is organized as follows. We give a brief review for the quadratic functors and the simplicial resolutions in Section 2. The determination of $\pi_4(\Sigma K(A,1))$ is given in section 3, where the proofs of Theorems 1.1 and 1.2 are also given. In section 4, we give case-by-case computations for $\pi_5(\Sigma K(A,1))$. In Section 5, we give some relations to K-theory.

2. THE QUADRATIC FUNCTORS AND THE SIMPLICIAL RESOLUTIONS

2.1. **Whitehead Quadratic Functor.** In [19, Chapter II], J. H. C Whitehead introduce the *universal quadratic functor* Γ_2 from abelian groups to abelian groups as follows: Let A be any abelian group. Then $\Gamma_2(A)$ is the group generated by the symbols $\gamma(x)$, one for each $x\in A$, subject to the defining relations

(1) $\gamma(-x)=\gamma(x)$;
(2) $\gamma(x+y+z)-\gamma(x+y)-\gamma(y+z)-\gamma(x+z)+\gamma(x)+\gamma(y)+\gamma(z)=0.$

Note. According to [19, p. 61], the group $\Gamma_2(A)$ is abelian and so the multiplication in $\Gamma_2(A)$ is denoted by $+$. Define

$$\gamma(x,y)=\gamma(x+y)-\gamma(x)-\gamma(y).$$

The following proposition helps for determining the group $\Gamma_2(A)$.

Proposition 2.1. [19, Theorem 5] *Let A be an abelian group with a basis $\{a_i \mid i\in I\}$ for a well-ordered index set I, and the defining relations $\{b_\lambda\equiv 0\}$. Then the group $\Gamma_2(A)$ is combinatorial defined by the set of symbolic generators $\gamma(a_i)$, $i\in I$, and $\gamma(a_i,a_j)$, $i,j\in I$ with $i<j$ with defining relations $\gamma(b_\lambda)\equiv 0$ and $\gamma(a_i,b_\lambda)\equiv 0$.*

Example 2.1. We list some examples of the group $\Gamma_2(A)$. The first two examples are direct consequences of the above proposition.

1. Let A be a free abelian group with a basis $\{a_i \mid i \in I\}$ for a well-ordered index set I. Then $\Gamma_2(A)$ i s the free abelian group with a basis given by $\gamma(a_i)$, $i \in I$, and $\gamma(a_i, a_j)$, $i, j \in I$ with $i < j$.
2. If A is a cyclic group of finite order m generated by a_1, then $\Gamma_2(A)$ is cyclic of order m or $2m$, according as m is odd or even, generated by $\gamma(a_1)$.
3. Let $A = \bigoplus_{i\in I} A_i$ for a well-ordered index set I. Then [19, Theorem 7]
$$\Gamma_2(A) \cong \bigoplus_{i\in I} \Gamma_2(A_i) \oplus \bigoplus_{\substack{i,j\in I\\ i<j}} A_i \otimes A_j.$$
4. For a general abelian group A, there is a short exact sequence [10, formula (13.8), p.93]
$$A \otimes A \overset{t}{\hookrightarrow} \Gamma_2(A) \twoheadrightarrow A \otimes \mathbb{Z}/2,$$
where $t(a \otimes b) = \gamma(a, b) = \gamma(a + b) - \gamma(a) - \gamma(b)$. □

2.2. **Lower Homology of $K(A, 2)$.** The homology of Eilenberg-MacLane spaces $K(A, n)$ has been studied in the classical reference [10] and other papers. See also [5] for the functorial description of homology groups of $K(A, 2)$ in all dimensions.

Lemma 2.1. [10, Theorems 20.5 and 21.1] *Let A be any abelian group. Then*

1. $H_2(K(A, 2)) = A$;
2. $H_3(K(A, 2)) = 0$;
3. $H_4(K(A, 2)) = \Gamma_2(A)$. □

The homology $H_5(K(A, 2))$ becomes a special functor on A. Let $R_2(A) = H_5(K(A, 2))$. The group $R_2(A)$ for finitely generated abelian group A can be computed as follows [10, Section 22]:

1) If A is a cyclic group of order infinite or odd, then $R_2(A) = 0$;
2) If $A = \mathbb{Z}/2^r\mathbb{Z}$ with $r \geq 1$, then $R_2(A) \cong \mathbb{Z}/2$.
3) Let $A = A_1 \oplus A_2$. Then $K(A, 2) \simeq K(A_1, 2) \times K(A_2, 2)$. By using Künneth theorem together with the fact that $H_1(K(A, 2)) = H_3(K(A, 2)) = 0$ from Lemma 2.1, we have
$$H_5(K(A, 2)) \cong H_5(K(A_1, 2)) \oplus H_5(K(A_2, 2)) \oplus \mathrm{Tor}(H_2(K(A_1, 2)), H_2(K(A_2, 2))).$$
Thus
$$R_2(A_1 \oplus A_2) \cong R_2(A_1) \oplus R_2(A_2) \oplus \mathrm{Tor}(A_1, A_2). \tag{2.1}$$

Recall the definition of the derived functors in the sense of Dold-Puppe [9]. Let F be an endofunctor in the category of abelian groups and A an abelian group. Take a projective resolution $P_* \to A$. Let N^{-1} be the inverse map to the normalization map due to Dold-Kan. Then $N^{-1}P_*$ is a free simplicial resolution of A. Then, the i-th derived functor of F applied to the abelian group A, is defined as follows:
$$L_iF(A) = \pi_i(F(N^{-1}P_*)),\ i \geq 0.$$

It is a well-known fact that this definition does not depend on a choice of a projective resolution. In these notations, one has a natural isomorphism:

$$R_2(A) = L_1\Gamma_2(A).$$

2.3. **Whitehead exact sequence.** Let X be a $(r-1)$-connected CW-complex, $r \geq 2$. There is the following long exact sequence of abelian groups [19, Theorem 1]:

$$\dots \to H_{n+1}(X) \to \Gamma_n(X) \to \pi_n(X) \xrightarrow{h_n} H_n(X) \to \Gamma_{n-1}(X) \to \dots, \tag{2.2}$$

where $\Gamma_n(X) = \mathrm{Im}(\pi_n(\mathrm{sk}_{n-1}(X)) \to \pi_n(\mathrm{sk}_n(X)))$ (here $\mathrm{sk}_i(X)$ is the i-th skeleton of X), h_n is the nth Hurewicz homomorphism.

The Hurewicz theorem is equivalent to the statement $\Gamma_i(X) = 0$, $i \leq r$. J. H. C. Whitehead computed the term $\Gamma_{r+1}(X)$: In the following theorem, assertion (1) was given in [19, Theorem 14] and assertion (2) was given the earlier paper [18]. According to the remarks in the end of [19, Section 14], assertion (2) has been discussed by G. W. Whitehead [20] as well.

Theorem 2.1. *Let X be a $(r-1)$-connected CW-complex with $r \geq 2$. Then*

1. *If $r = 2$, then $\Gamma_3(X) \cong \Gamma_2(\pi_2(X))$.*
2. *If $r > 2$, then $\Gamma_{r+1}(X) \cong \pi_r(X) \otimes \mathbb{Z}/2$.* □

The isomorphism $\Gamma_2(\pi_2(X)) \to \Gamma_3(X)$ is constructed as follows: Let $\eta\colon S^3 \to S^2$ be the Hopf map and let $x \in \pi_2(X)$ be written as the the composite

$$S^2 \xrightarrow{\tilde{x}} \mathrm{sk}_2(X) \hookrightarrow \mathrm{sk}_3(X).$$

Then the composite

$$S^3 \xrightarrow{\eta} S^2 \xrightarrow{\tilde{x}} \mathrm{sk}_2(X) \hookrightarrow \mathrm{sk}_3(X)$$

defines an element $\eta^*(x) \in \Gamma_3(X)$. According to [19, Section 13], the mapping

$$\eta_1\colon \Gamma_2(\pi_2(X)) \to \Gamma_3(X), \quad \gamma(x) \mapsto \eta^*(x), \tag{2.3}$$

is a well-defined isomorphism of groups. The construction of the isomorphism $\pi_r(X) \otimes \mathbb{Z}/2 \to \Gamma_{r+1}(X)$ in assertion (2) is similar.

Recall the description of the functors $\Gamma_{r+2}(X)$ due to H.-J. Baues [2]. Consider the third super-Lie functor

$$\mathcal{L}_s^3 : \mathsf{Ab} \to \mathsf{Ab}$$

defined as

$$\mathcal{L}_s^3(A) = im\{A \otimes A \otimes A \xrightarrow{l} A \otimes A \otimes A\}$$

where

$$l(a \otimes b \otimes c) = \{a, b, c\} := a \otimes b \otimes c + b \otimes a \otimes c - c \otimes a \otimes b - c \otimes b \otimes a, \ a, b, c \in A.$$

Observe that $\mathcal{L}_s^3(A) = \ker\{A \otimes \Lambda^2(A) \xrightarrow{r} \Lambda^3(A)\}$, where $\Lambda^i(A)$ is the ith exterior power of A and the map r is given as

$$r(a \otimes b \wedge c) = a \wedge b \wedge c, \ a, b, c \in A.$$

Let the complex X be simply connected. Given an abelian group A, define the map

$$q : \Gamma_2(A) \otimes A \to \mathcal{L}_s^3(A) \oplus \Gamma_2(A) \otimes \mathbb{Z}/2$$

by setting

$$q(\gamma_2(a) \otimes b) = -\{b, a, a\} + (\gamma_2(a+b) - \gamma_2(a) - \gamma_2(b)) \otimes 1, \ a, b \in A.$$

Define the group $\Gamma_2^2 X = \Gamma_2^2(\Gamma_2(\pi_2 X) \to \pi_3 X)$ as the pushout:

$$\begin{tikzcd}
\Gamma_2(\pi_2(X)) \otimes (\pi_2(X) \oplus \mathbb{Z}/2) \arrow[r, "q\oplus id"] \arrow[d, "\eta_1\otimes id"] & \mathcal{L}_s^3(\pi_2(X)) \oplus \Gamma_2(\pi_2(X)) \otimes \mathbb{Z}/2 \arrow[d] \\
\pi_3(X) \otimes (\pi_2(X) \oplus \mathbb{Z}/2) \arrow[r] & \Gamma_2^2(X)
\end{tikzcd} \tag{2.4}$$

Theorem 2.2. [2, Theorem 3.1] *Let X be a $(r-1)$-connected CW-complexes with $r \geq 2$.*

1) *If $r = 2$, then there is a natural short exact sequence*

$$0 \to \Gamma_2^2(X) \to \Gamma_4(X) \to R_2(\pi_2(X)) \to 0.$$

2) *If $r = 3$, then there is a natural exact sequence*

$$0 \to \pi_4(X) \otimes \mathbb{Z}/2 \oplus \Lambda^2(\pi_3(X)) \to \Gamma_5(X) \to \mathrm{Tor}(\pi_3(X), \mathbb{Z}/2) \to 0.$$

3) *If $r \geq 4$, there is a natural exact sequence*

$$0 \to \pi_{r+1}(X) \otimes \mathbb{Z}/2 \to \Gamma_{r+2}(X) \to \mathrm{Tor}(\pi_r(X), \mathbb{Z}/2) \to 0.$$

□

Let A be an abelian group. Consider the Hurewicz homomorphism

$$h_*\colon \pi_*(\Sigma K(A,1)) \to \tilde{H}_*(\Sigma K(A,1)) = \tilde{H}_{*-1}(K(A,1)) = \tilde{H}_{*-1}(A).$$

Since $H_*(K(A,1))$ is graded commutative ring, the inclusion

$$A = H_1(K(A,1)) \hookrightarrow H_*(K(A,1))$$

induces a ring homomorphism

$$\lambda\colon \Lambda(A) \longrightarrow H_*(K(A,1)).$$

By [10, Theorem 19.3], λ is a monomorphism and so we may consider $\Lambda^n(A) \subseteq H_n(K(A,1)) = H_n(A)$.

Lemma 2.2. *For every abelian group A, the Hurewicz image*

$$\mathrm{Im}(h_{n+1}\colon \pi_{n+1}(\Sigma K(A,1)) \to H_n(A))$$

contains the subgroup $\Lambda^n(A)$.

Proof. From the naturality, it suffices to show that the statement holds for a free abelian group A.

When A is a free abelian group, then $K(A,1)$ is a (weak) Cartesian product of the circles. Thus $\Sigma K(A,1)$ is a wedge of spheres from the suspension splitting that

$$\Sigma X \times Y \simeq \Sigma X \vee \Sigma Y \vee \Sigma X \wedge Y$$

and so the Hurewicz homomorphism induces an epimorphism

$$h_*\colon \pi_{n+1}(\Sigma K(A,1)) \twoheadrightarrow H_n(A) = \Lambda^n(A)$$

for a free abelian group A. This finishes the proof. □

2.4. **Carlsson construction.** Let G_* be a simplicial group and X a pointed simplicial set with a base point $*$. Consider the simplicial group $F^{G_*}(X)$ defined as

$$F^H(X)_n = \coprod_{x \in X_n} (G_n)_x,$$

i.e. in each degree $F^{G_*}(X)_n$ is the free product of groups G_n numerated by elements of X_n modulo $(G_n)_*$, with the canonical choice of face and degeneracy morphisms. It is proved in [7] that the geometric realization $|F^{G_*}(X)|$ is homotopy equivalent to the loop space $\Omega(|X| \wedge B|G|)$. The main example we will consider is the simplicial circle $X = S^1$ with

$$S^1_0 = \{*\},\ S^1_1 = \{*, \sigma\},\ S^1_2 = \{*, s_0\sigma, s_1\sigma\}, \dots, S^1_n = \{*, x_0, \dots, x_n\},$$

where $x_i = s_n \dots \hat{s}_i \dots s_0 \sigma$ and the simplicial group G_* with $G_n = G$ for a given group G, with identity homomorphisms as all face and degeneracy maps. In this case we use the notation $F^G(X) = F^{G_*}(X)$. One has a homotopy equivalence

$$|F^G(S^1)| \simeq \Omega\Sigma K(G,1).$$

The group $F^G(S^1)_n$ is the n-fold free product of G:

$$F^G(S^1)_1 = G,\ F^G(S^1)_2 = G * G,\ F^G(S^1)_3 = G * G * G,\ \dots$$

We can formally identify $G * G$ with $s_0 G * s_1 G$, $G * G * G$ with $s_1 s_0 G * s_2 s_0 G * s_2 s_1 G$, etc, and to define naturally the face and degeneracy maps:

$$\begin{tikzcd}
F^{G_*}(S^1): \quad \dots \arrow[r, shift left=4] \arrow[r, phantom, shift left=2, "\cdots"] \arrow[r, shift left=1] \arrow[r, leftarrow, shift right=1] \arrow[r, phantom, shift right=2, "\cdots"] \arrow[r, leftarrow, shift right=4] & G * G * G \arrow[r, shift left=7] \arrow[r, shift left=5] \arrow[r, shift left=3] \arrow[r, shift left=1] \arrow[r, leftarrow, shift right=1] \arrow[r, leftarrow, shift right=3] \arrow[r, leftarrow, shift right=5] & G * G \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[r, leftarrow, shift right=2] \arrow[r, leftarrow, shift right=4] & G.
\end{tikzcd}$$

Remark. Consider the second term $F^G(S^1)_2 = G * G$ and face morphisms $d_0, d_1, d_2 : G * G = s_0(G) * s_1(G) \to G$ defined as

$$d_0 : \begin{cases} s_0(g) \mapsto g \\ s_1(g) \mapsto 1 \end{cases},\ d_1 : \begin{cases} s_0(g) \mapsto g \\ s_1(g) \mapsto g \end{cases},\ d_2 : \begin{cases} s_0(g) \mapsto 1 \\ s_1(g) \mapsto g \end{cases}.$$

There is a natural commutative diagram

$$\begin{tikzcd}
\pi_3(\Sigma K(G,1)) \arrow[r, hook] \arrow[d, "\simeq"] & G \otimes G \arrow[r] \arrow[d, "f"] & G \arrow[r, two heads] \arrow[d, "\simeq"] & G_{ab} \arrow[d, "\simeq"] \\
\pi_3(\Sigma K(G,1)) \arrow[r, hook] & (ker(d_1) \cap ker(d_2))/\mathcal{B}_2 \arrow[r] & G \arrow[r, two heads] & G_{ab}
\end{tikzcd} \tag{2.5}$$

where $\mathcal{B}_2$ is the 2-boundary subgroup of $G * G$ and the map f is defined as[1]

$$f(g \otimes h) = [s_0(g)s_1(g)^{-1}, s_0(h)].\mathcal{B}_2.$$

There is a natural description of the 2-boundary (see [11], for example):

$$\mathcal{B}_2 = [ker(d_0), ker(d_1) \cap ker(d_2)][ker(d_1), ker(d_2) \cap ker(d_0)][ker(d_2), ker(d_0) \cap ker(d_1)].$$

Diagram (2.5) implies that f is a natural isomorphism.

---

[1] We use the standard commutator relations: $[g,h] = g^{-1}h^{-1}gh$

In the case $G = \mathbb{Z}$, the simplicial group $F^G(S^1)$ is identical to the Milnor construction $F(S^1)$, with $F(S^1)_n$ a free group of rank n, for $n \geq 1$:

$$\begin{tikzcd}
F(S^1): \quad \dots \arrow[r, shift left=4] \arrow[r, phantom, shift left=2, "\cdots"] \arrow[r, leftarrow] \arrow[r, phantom, shift right=2, "\cdots"] \arrow[r, leftarrow, shift right=4] & F_3 \arrow[r, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[r, leftarrow, shift right=2] \arrow[r, leftarrow, shift right=4] \arrow[r, leftarrow, shift right=6] & F_2 \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[r, leftarrow, shift right=2] \arrow[r, leftarrow, shift right=4] & \mathbb{Z}.
\end{tikzcd}$$

In this case there is a homotopy equivalence

$$|F(S^1)| \simeq \Omega S^2$$

and the construction $F(S^1)$ provides a combinatorial model for the computation of homotopy groups of the 2-sphere S^2. The construction $F(S^1)$ was studied from the group-theoretical point of view in [21]. It is easy to find the simplicial generators of the homotopy classes of $\pi_i(F(S^1)) = \pi_{i+1}(S^2)$ for $i = 3, 4, 5$. In order to find these simplicial generators, consider the sequence of maps between Milnor simplicial constructions $F(S^4) \to F(S^3) \to F(S^2) \to F(S^1)$ such that the induced homomorphisms $\mathbb{Z} = \pi_2(F(S^2)) \to \pi_2(F(S^1)) = \mathbb{Z}$ and $\mathbb{Z} = \pi_3(F(S^3)) \to \pi_3(F(S^2)) = \mathbb{Z}/2$ are epimorphisms and define the homotopy classes of $\pi_3(S^2)$ and $\pi_4(S^3)$ respectively.

$$\begin{tikzcd}
F(S^3)_4 \arrow[r, shift left=4] \arrow[r, phantom, shift left=2, "\cdots"] \arrow[r, leftarrow] \arrow[r, phantom, shift right=2, "\cdots"] \arrow[r, leftarrow, shift right=4] \arrow[d] & \mathbb{Z} \arrow[d, "\eta^2"] & & \\
F(S^2)_4 \arrow[r, shift left=4] \arrow[r, phantom, shift left=2, "\cdots"] \arrow[r, leftarrow] \arrow[r, phantom, shift right=2, "\cdots"] \arrow[r, leftarrow, shift right=4] \arrow[d] & F(S^2)_3 \arrow[r, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[r, leftarrow, shift right=2] \arrow[r, leftarrow, shift right=4] \arrow[r, leftarrow, shift right=6] \arrow[d] & \mathbb{Z} \arrow[d, "\eta"] & \\
F(S^1)_4 \arrow[r, shift left=4] \arrow[r, phantom, shift left=2, "\cdots"] \arrow[r, leftarrow] \arrow[r, phantom, shift right=2, "\cdots"] \arrow[r, leftarrow, shift right=4] & F(S^1)_3 \arrow[r, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[r, leftarrow, shift right=2] \arrow[r, leftarrow, shift right=4] \arrow[r, leftarrow, shift right=6] & F(S^1)_2 \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[r, leftarrow, shift right=2] \arrow[r, leftarrow, shift right=4] & \mathbb{Z}
\end{tikzcd}$$

For $n \geq 3$, the homotopy class of $\pi_n(S^{n-1})$ defined as $\pi_{n-1}(F(S^{n-2}))$ is generated by $[s_0(\sigma_{n-2}), s_1(\sigma_{n-2})]$ in $F(S^{n-2})_{n-1}$ (see [21]), where σ_{n-2} is a generator of $F(S^{n-2})_{n-2} = \mathbb{Z}$. That is, we can define the simplicial suspension maps $\eta^i : F(S^{i+1})_{i+1} \to F(S^i)_{i+1}$ by

$$\eta^i : \sigma_{i+1} \to [s_0(\sigma_i), s_1(\sigma_i)], \ i \geq 1.$$

Since the generators of $\pi_i(S^2)$ are presented by suspensions over Hopf fibration for $i = 3, 4, 5$, the simplicial generators of $\pi_i(F(S^1))$, $i = 2, 3, 4$ are given by the following elements:

$$w_2(x_0, x_1) = [x_0, x_1] \tag{2.6}$$

$$w_3(x_0, x_1, x_2) = [[x_0, x_2], [x_0, x_1]] \tag{2.7}$$

$$w_4(x_0, x_1, x_2, x_3) = [[[x_0, x_3], [x_0, x_1]], [[x_0, x_2], [x_0, x_1]]]. \tag{2.8}$$

Here we use the natural notations $x_j := s_i \dots \hat{s}_j \dots s_0(\sigma_1)$, $j = 0, \dots, i$ for the basis elements in $F(S^1)_{i+1}$[2].

---

[2] One can continue the process of construction of elements $w_{n+1}(x_0, \dots, x_n)$ by the following law: $w_{n+1}(x_0, \dots, x_n) = [w_n(x_0, \dots, \hat{x}_{n-1}, x_n), w_n(x_0, \dots, x_{n-1})]$. In this case, the 16-commutator bracket $w_5(x_0, \dots, x_4)$ corresponds to the element of order 2 in $\pi_6(S^2)$, but the 32-commutator bracket

2.5. **Spectral sequence.** Consider an abelian group A and its two-step flat resolution

$$0 \to A_1 \to A_0 \to A \to 0.$$

By Dold-Kan correspondence, we obtain the following free abelian simplicial resolution of A:

$$\begin{tikzcd}
N^{-1}(A_1 \hookrightarrow A_0): \quad \cdots \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] & A_1 \oplus s_0(A_0) \arrow[l, shift left=2] \arrow[l, shift left=4] \arrow[r, shift left=2] \arrow[r] & A_0. \arrow[l, shift left=2]
\end{tikzcd}$$

Applying Carlsson construction to the resolution $N^{-1}(A_1 \hookrightarrow A_0)$, we obtain the following bisimplicial group:

$$\begin{tikzcd}
F^{N^{-1}(A_1\hookrightarrow A_0)_2}(S^n)_3 \arrow[r, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[d, shift left=4] \arrow[d, shift left=2] \arrow[d] & F^{N^{-1}(A_1\hookrightarrow A_0)_2}(S^n)_2 \arrow[l, shift left=2] \arrow[l, shift left=4] \arrow[l, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[d, shift left=4] \arrow[d, shift left=2] \arrow[d] & N^{-1}(A_1 \hookrightarrow A_0)_2 \arrow[l, shift left=2] \arrow[l, shift left=4] \arrow[d, shift left=4] \arrow[d, shift left=2] \arrow[d] \\
F^{A_1\oplus s_0(A_0)}(S^n)_3 \arrow[u, shift left=2] \arrow[u, shift left=4] \arrow[r, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[d, shift left=2] \arrow[d] & F^{A_1\oplus s_0(A_0)}(S^n)_2 \arrow[u, shift left=2] \arrow[u, shift left=4] \arrow[l, shift left=2] \arrow[l, shift left=4] \arrow[l, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[d, shift left=2] \arrow[d] & A_1 \oplus s_0(A_0) \arrow[u, shift left=2] \arrow[u, shift left=4] \arrow[l, shift left=2] \arrow[l, shift left=4] \arrow[d, shift left=2] \arrow[d] \\
F^{A_0}(S^n)_3 \arrow[u, shift left=2] \arrow[r, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] & F^{A_0}(S^n)_2 \arrow[u, shift left=2] \arrow[l, shift left=2] \arrow[l, shift left=4] \arrow[l, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] & A_0 \arrow[u, shift left=2] \arrow[l, shift left=2] \arrow[l, shift left=4]
\end{tikzcd}$$

Here the mth horizontal simplicial group is Carlsson construction $F^{N^{-1}(A_1\hookrightarrow A_0)_m}(S^n)$. By the result of Quillen [15], we obtain the following spectral sequence:

$$E^2_{p,q} = \pi_q(\pi_p(\Sigma^n K(N^{-1}(A_1 \hookrightarrow A_0), 1)) \Longrightarrow \pi_{p+q}(\Sigma^n K(A, 1)). \tag{2.9}$$

Consider now a non-abelian analog of this spectral sequence, for $n = 1$. Suppose now that a group G is arbitrary, not necessary abelian. Consider a simplicial resolution of G:

$$G_\bullet \to G,$$

i.e. $G_\bullet$ is a simplicial group with $\pi_0(G_\bullet) = G,\ \pi_i(G_\bullet) = 0,\ i > 0$. Consider the following bisimplicial group

$$\begin{tikzcd}
G_2 * G_2 * G_2 \arrow[r, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[d, shift left=4] \arrow[d, shift left=2] \arrow[d] & G_2 * G_2 \arrow[l, shift left=2] \arrow[l, shift left=4] \arrow[l, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[d, shift left=4] \arrow[d, shift left=2] \arrow[d] & G_2 \arrow[l, shift left=2] \arrow[l, shift left=4] \arrow[d, shift left=4] \arrow[d, shift left=2] \arrow[d] \\
G_1 * G_1 * G_1 \arrow[u, shift left=2] \arrow[u, shift left=4] \arrow[r, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[d, shift left=2] \arrow[d] & G_1 * G_1 \arrow[u, shift left=2] \arrow[u, shift left=4] \arrow[l, shift left=2] \arrow[l, shift left=4] \arrow[l, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] \arrow[d, shift left=2] \arrow[d] & G_1 \arrow[u, shift left=2] \arrow[u, shift left=4] \arrow[l, shift left=2] \arrow[l, shift left=4] \arrow[d, shift left=2] \arrow[d] \\
G_0 * G_0 * G_0 \arrow[u, shift left=2] \arrow[r, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] & G_0 * G_0 \arrow[u, shift left=2] \arrow[l, shift left=2] \arrow[l, shift left=4] \arrow[l, shift left=6] \arrow[r, shift left=4] \arrow[r, shift left=2] \arrow[r] & G_0 \arrow[u, shift left=2] \arrow[l, shift left=2] \arrow[l, shift left=4]
\end{tikzcd}$$

Again, by the result of Quillen [15], we obtain the following spectral sequence:

$$E^2_{p,q} = \pi_q(\pi_p(\Sigma K(G_\bullet, 1))) \Longrightarrow \pi_{p+q}(\Sigma K(G, 1)). \tag{2.10}$$

$w_6(x_0, \dots, x_5)$ lies in the simplicial boundary subgroup $\mathcal{B}F(S^1)_6$ (see [10]). The construction of a simplicial generator of the 3-torsion in $\pi_6(S^2)$ is more tricky: it is possible to find its simplicial representative which is a product of six brackets of the commutator weight six.

If $G_\bullet$ is a free simplicial resolution, the spectral sequence (2.10) contains a lot of canonical differentials of a complicated nature:

$$\begin{tikzcd}
 & & H_4(G) \arrow[ddl] \\
\pi_2(\mathcal{L}_s^3((G_\bullet)_{ab}) \oplus \Gamma_2((G_\bullet)_{ab}) \otimes \mathbb{Z}/2) & \pi_2(\Gamma_2((G_\bullet)_{ab})) & H_3(G) \arrow[ddl] \\
\pi_1(\mathcal{L}_s^3((G_\bullet)_{ab}) \oplus \Gamma_2((G_\bullet)_{ab}) \otimes \mathbb{Z}/2) & \pi_1(\Gamma_2((G_\bullet)_{ab})) & H_2(G) \\
\mathcal{L}_s^3(G_{ab}) \oplus \Gamma_2(G_{ab}) \otimes \mathbb{Z}/2 & \Gamma_2(G_{ab}) & G_{ab}
\end{tikzcd}$$

3. On group $\pi_4(\Sigma K(A,1))$

3.1. **The group $\pi_4(\Sigma K(A,1))$ for an abelian group A.** Let A be an abelian group. Consider the homotopy commutative diagram of fibre sequences

$$\begin{tikzcd}
\Sigma K(A,1) \wedge K(A,1) \arrow[r, "H"] \arrow[d, equal] & \Sigma K(A,1) \arrow[r] \arrow[d] \arrow[dr, phantom, "\text{pull}"] & K(A,2) = BK(A,1) \arrow[d, "\Delta"] \\
\Sigma K(A,1) \wedge K(A,1) \arrow[r, "f"] & K(A,2) \vee K(A,2) \arrow[r, hook] & K(A,2) \times K(A,2),
\end{tikzcd} \tag{3.1}$$

where H is the Hopf fibration. Thus we have the following lemma:

Lemma 3.1. *There are isomorphisms*

$$\pi_n(\Sigma K(A,1) \wedge K(A,1)) \cong \pi_n(\Sigma K(A,1)) \cong \pi_n(K(A,2) \vee K(A,2))$$

for $n \geq 3$. In particular, $\pi_3(\Sigma K(A,1)) \cong \pi_3(\Sigma K(A,1) \wedge K(A,1)) \cong A \otimes A$. □

By Lemma 2.1, the lower homology of the wedge $K(A,2) \vee K(A,2)$ are the following:

Lemma 3.2.

$$\begin{aligned}
&H_2(K(A,2) \vee K(A,2)) = A \oplus A,\\
&H_3(K(A,2) \vee K(A,2)) = 0,\\
&H_4(K(A,2) \vee K(A,2)) = \Gamma_2(A) \oplus \Gamma_2(A),\\
&H_5(K(A,2) \vee K(A,2)) = R_2(A) \oplus R_2(A).
\end{aligned}$$

□

Lemma 3.3. *The Hurewicz image*

$$h_n \colon \pi_n(K(A,2) \vee K(A,2)) \longrightarrow H_n(K(A,2) \vee K(A,2))$$

is zero for $n \geq 3$.

Proof. The assertion follows from the commutative diagram

$$\begin{tikzcd}
\pi_n(K(A,2)\vee K(A,2)) \arrow[r, "h_n"] \arrow[d] & H_n(K(A,2)\vee K(A,2)) \arrow[d, hook] \\
\pi_n(K(A,2)\times K(A,2))=0 \arrow[r, "h_n"] & H_n(K(A,2)\times K(A,2)).
\end{tikzcd}$$

□

Theorem 3.1. *Let A be any abelian group. Then there is a natural short exact sequence*

$$(\Lambda^2(A)\otimes A)^{\oplus 2}\oplus A\otimes A\otimes\mathbb{Z}/2 \hookrightarrow \pi_4(\Sigma K(A,1)) \twoheadrightarrow \mathrm{Tor}(A,A).$$

Moreover $(\Lambda^2(A)\otimes A)^{\oplus 2}$ *is an (unnatural) summand of* $\pi_4(\Sigma K(A,1))$.

Proof. Let $X=K(A,2)\vee K(A,2)$ and let $Y=K(A,2)\times K(A,2)=K(A\oplus A,2)$. From Lemmas 3.2 and 3.3, there is a short exact sequence

$$R_2(A)\oplus R_2(A)\hookrightarrow \Gamma_4(X)\twoheadrightarrow \pi_4(X).$$

The inclusion $j\colon X=K(A,2)\vee K(A,2)\hookrightarrow Y=K(A,2)\times K(A,2)$ induces a commutative diagram

$$\begin{tikzcd}
H_5(X)=R_2(A)\oplus R_2(A) \arrow[r, hook] \arrow[d, hook, "j_*"] & \Gamma_4(X) \arrow[r, two heads] \arrow[d, "j_*"] & \pi_4(X) \arrow[d, "j_*"] \\
H_5(Y)=R_2(A\oplus A) \arrow[r, "\cong", "\phi"'] & \Gamma_4(Y) \arrow[r] & \pi_4(Y)=0.
\end{tikzcd} \tag{3.2}$$

By formula (2.1),

$$H_5(K(A,2)\times K(A,2))=H_5(K(A\oplus A,2))=R_2(A)\oplus R_2(A)\oplus\mathrm{Tor}(A,A)$$

and so the cokernel of $j_*\colon H_5(X)\to H_5(Y)$ is $\mathrm{Tor}(A,A)$. On the other hand, from Theorem 2.2(1), there is a commutative diagram of short exact sequences

$$\begin{tikzcd}
\Gamma_2^2(X) \arrow[r, hook] \arrow[d] & \Gamma_4(X) \arrow[r, two heads] \arrow[d, "j_*"] & R_2(\pi_2(X))=R_2(A\oplus A) \arrow[d, "\cong"] \\
\Gamma_2^2(Y) \arrow[r, hook] & \Gamma_4(Y) \arrow[r, two heads, "\psi"] & R_2(\pi_2(Y))=R_2(A\oplus A).
\end{tikzcd}$$

The composite

$$\psi\circ\phi\colon R_2(A\oplus A)\xrightarrow[\phi]{\cong}\Gamma_4(Y)\underset{\psi}{\twoheadrightarrow}R_2(A\oplus A)$$

is a natural self epimorphism for any abelian group A. It is an isomorphism for any finitely generated abelian A and so an isomorphism for any abelian group A by considering the direct limit. Thus $j_*\colon\Gamma_4(X)\to\Gamma_4(Y)$ is an epimorphism and, from diagram (3.2), there is a short exact sequence

$$\Gamma_2^2(X)\hookrightarrow\pi_4(X)\twoheadrightarrow\mathrm{Tor}(A,A). \tag{3.3}$$

Let $Z=\Sigma K(A,1)\wedge K(A,1)$ and let $f\colon Z\to X$ be the map in diagram (3.1). Consider the commutative diagram

$$\begin{tikzcd}
\pi_3(Z)\otimes\mathbb{Z}/2=\Gamma^2_2(Z) \arrow[r, hook, "\cong"] \arrow[d, "f_*"] & \Gamma_4(Z) \arrow[r] \arrow[d, "f_*"] & \pi_4(Z) \arrow[r] \arrow[d, "f_*"', "\cong"] & H_4(Z) \arrow[r] & \Gamma_3(Z)=0 \\
\Gamma^2_2(X) \arrow[r, hook] & \Gamma_4(X) \arrow[r, two heads] & \pi_4(X),
\end{tikzcd}\tag{3.4}$$

where $\Gamma^2_2(Z)\to\Gamma_4(Z)$ is an isomorphism because its cokernel $R_2(\pi_2(Z))=0$. From the definition (2.4) of the functor Γ^2_2,

$$f_*\colon\Gamma^2_2(Z)\longrightarrow\Gamma^2_2(X)$$

is a monomorphism with retracting homomorphism $\phi'\colon\Gamma^2_2(X)\to\Gamma^2_2(Z)$. By the short exact sequence (3.3), $\Gamma_4(Z)\to\pi_4(Z)$ is a monomorphism and so there is a short exact sequence

$$A\otimes A\otimes\mathbb{Z}/2=\Gamma_4(Z)\hookrightarrow\pi_4(Z)\twoheadrightarrow H_4(Z)=H_3(K(A,1)\wedge K(A,1)).\tag{3.5}$$

Note that $H_1(K(A,1))=A$ and $H_2(K(A,1))=\Lambda^2(A)$. By the Künneth theorem, there is a natural short exact sequence

$$(A\otimes\Lambda^2(A))^{\oplus 2}\hookrightarrow H_4(Z)\overset{\psi'}{\twoheadrightarrow}\mathrm{Tor}(A,A).$$

Consider the composite

$$\theta_A\colon\Gamma^2_2(X)=\Gamma^2_2(\Gamma_2(A\oplus A)\to A\otimes A)\hookrightarrow\pi_4(X)\xrightarrow{f_*^{-1}}\pi_4(Z)\twoheadrightarrow H_4(Z)\overset{\psi'}{\twoheadrightarrow}\mathrm{Tor}(A,A),$$

which is natural on any abelian group A. If A is a free abelian group, then $\theta_A=0$. For any abelian group A, choose any free abelian group A_0 with an epimorphism $g\colon A_0\twoheadrightarrow A$. From the definition (2.4) of Γ^2_2,

$$\Gamma^2_2(g)\colon\Gamma^2_2(\Gamma_2(A_0\oplus A_0)\to A_0\otimes A_0)\longrightarrow\Gamma^2_2(\Gamma_2(A\oplus A)\to A\otimes A)$$

is an epimorphism. By the naturality of θ_A, we have $\theta_A=0$ because $\theta_{A_0}=0$. Now, from diagram (3.4), there is a commutative diagram of natural short exact sequences

$$\begin{tikzcd}
A\otimes A\otimes\mathbb{Z}/2 \arrow[r, equal] \arrow[d, hook, "\circlearrowleft"] & A\otimes A\otimes\mathbb{Z}/2 \arrow[d, hook] & \\
\Gamma^2_2(X) \arrow[r, hook] \arrow[d] & \pi_4(\Sigma K(A,1)) \arrow[r, two heads] \arrow[d] & \mathrm{Tor}(A,A) \arrow[d, "\cong"] \\
(A\otimes\Lambda^2(A))^{\oplus 2} \arrow[r, hook] & H_4(Z) \arrow[r, two heads] & \mathrm{Tor}(A,A).
\end{tikzcd}$$

It follows that there is a natural (on A) isomorphism

$$\Gamma^2_2(X)\cong A\otimes A\otimes\mathbb{Z}/2\oplus(A\otimes\Lambda^2(A))^{\oplus 2}.$$

Since $(A\otimes\Lambda^2(A))^{\oplus 2}$ is an (unnatural) summand of $H_4(Z)$, it is an (unnatural) summand of $\pi_4(\Sigma K(A,1))$. The proof is finished. □

Corollary 3.1. *Let p be an odd prime integer. Then*

$$\pi_4(\Sigma K(\mathbb{Z}/p^r,1))=\mathbb{Z}/p^r$$

and the Hurewicz homomorphism

$$\pi_4(\Sigma K(\mathbb{Z}/p^r,1))\to H_4(\Sigma K(\mathbb{Z}/p^r,1))$$

is an isomorphism.

Proof. In this case, $A\otimes A\otimes\mathbb{Z}/2=0$. Since $\mathbb{Z}/p^r$ is cyclic, $\Lambda^2(A)\otimes A=0$ and hence the result. □

For completely determining the group $\pi_4(\Sigma K(A,1))$, we have to consider the divisibility problem of the elements in the subgroup $A\otimes A\otimes\mathbb{Z}/2=\Gamma_4(Z)\subseteq\pi_4(\Sigma K(A,1))=\pi_4(Z)$. We solve this problem for any finitely generated abelian group A.

Lemma 3.4. *Let A be any abelian group and let $j\colon M(A,1)\to K(A,1)$ be a map such that $j_*\colon H_1(M(A,1))\to H_1(K(A,1))$ is an isomorphism. Then there is an (unnatural) splitting exact sequence*

$$\begin{tikzcd}\pi_4(\Sigma M(A,1)\wedge M(A,1)) \arrow[r, hook, "\curvearrowleft", "(\Sigma j\wedge j)_*"'] & \pi_4(\Sigma K(A,1)\wedge K(A,1)) \arrow[r, two heads, "\curvearrowleft"] & (A\otimes\Lambda^2(A))^{\oplus 2}.\end{tikzcd}$$

Proof. Let $X=\Sigma M(A,1)\wedge M(A,1)$ and let $Z=\Sigma K(A,1)\wedge K(A,1)$. The assertion follows from the commutative diagram of short exact sequences

$$\begin{tikzcd}\Gamma_4(X) \arrow[r, hook] \arrow[d, "\cong"] & \pi_4(X) \arrow[r, two heads] \arrow[d] & H_4(X)=\mathrm{Tor}(A,A) \arrow[d, hook] \\ \Gamma_4(Z) \arrow[r, hook] & \pi_4(Z) \arrow[r, two heads] & H_4(Z),\end{tikzcd}$$

where the bottom row is short exact by equation (3.5). □

Given a finitely generated abelian group A, let

$$A=A_1\oplus\bigoplus_{\substack{r\geq 1\\ p\text{ is a prime}}}A_{p^r}$$

be the primary decomposition of A, where A_1 is torsion free and A_{p^r} is a free $\mathbb{Z}/p^r$-module.

Theorem 3.2. *Let A be any finitely generated abelian group. Let $A=A_2\oplus B$ with $B=A_1\oplus\bigoplus_{p^r\neq 2}A_{p^r}$. Then*

$$\begin{aligned}\pi_4(\Sigma K(A,1)) &\cong \tfrac{1}{2}(A_2\otimes A_2)\oplus(A_2\otimes B)^{\oplus 2}\oplus B^{\otimes 2}\otimes\mathbb{Z}/2\oplus(A\otimes\Lambda^2(A))^{\oplus 2}\\ &\quad\oplus\mathrm{Tor}(A_2,B)^{\oplus 2}\oplus\mathrm{Tor}(B,B),\end{aligned}$$

where $\frac{1}{2}(A_2\otimes A_2)$ is a free $\mathbb{Z}/4$-module with rank of $\dim_{\mathbb{Z}/2}(A_2\otimes A_2)$.

Proof. Let $X=\Sigma M(A,1)\wedge M(A,1)$. By Lemma 3.4, it suffices to show that

$$\pi_4(X)\cong\frac{1}{2}(A_2\otimes A_2)\oplus(A_2\otimes B)^{\oplus 2}\oplus B^{\otimes 2}\otimes\mathbb{Z}/2\oplus\mathrm{Tor}(A_2,B)^{\oplus 2}\oplus\mathrm{Tor}(B,B).$$

Observe that there is a homotopy decomposition

$$X\simeq\bigvee_{\substack{r,s\geq 0\\ p,q\text{ prime}}}\Sigma M(A_{p^r},1)\wedge M(A_{q^s},1),$$

where we allow r,s to be 0 for having the factor A_1 to be appeared. Thus there is a decomposition

$$\pi_4(X)\cong\bigoplus_{\substack{r,s\geq 0\\ p,q\text{ prime}}}\pi_4(\Sigma M(A_{p^r},1)\wedge M(A_{q^s},1)).\tag{3.6}$$

Let $r,s\geq 1$ and let p and q be positive prime integers. From [13, Corollary 6.6], there is a homotopy decomposition

$$\begin{aligned}&\Sigma M(\mathbb{Z}/p^r,1)\wedge M(\mathbb{Z}/q^s,1)\\ &\simeq\begin{cases}* & \text{if}\quad p\neq q,\\ M(\mathbb{Z}/p^{\min\{r,s\}},3)\vee M(\mathbb{Z}/p^{\min\{r,s\}},4) & \text{if}\quad p=q\text{ and }\max\{p^r,q^s\}>2.\end{cases}\end{aligned}\tag{3.7}$$

By taking π_4 to above decomposition, we have

$$\pi_4(\Sigma M(\mathbb{Z}/p^r,1)\wedge M(\mathbb{Z}/q^s,1))\cong\mathbb{Z}/p^r\otimes\mathbb{Z}/q^s\otimes\mathbb{Z}/2\oplus\mathrm{Tor}(\mathbb{Z}/2^r,\mathbb{Z}/2^s)\tag{3.8}$$

if $\max\{p^r,q^s\}>2$. Clearly this formula also holds for the case where $p^r=1$ or $q^s=1$. For the case $p^r=q^s=2$, we claim that

$$\pi_4(\Sigma M(\mathbb{Z}/2,1)\wedge M(\mathbb{Z}/2,1)))=\mathbb{Z}/4.\tag{3.9}$$

Let $Y=\Sigma M(\mathbb{Z}/2,1)\wedge M(\mathbb{Z}/2,1)$. From the short exact sequence

$$\Gamma_4(Y)=\mathbb{Z}/2\hookrightarrow\pi_4(Y)\twoheadrightarrow H_4(Y)=\mathbb{Z}/2,$$

the group $\pi_4(Y)=\mathbb{Z}/4$ or $\mathbb{Z}/2\oplus\mathbb{Z}/2$. Suppose that $\pi_4(Y)=\mathbb{Z}/2\oplus\mathbb{Z}/2$. Then there exists an element $\alpha\in\pi_4(Y)$ of order 2 which has the nontrivial Hurewicz image. Since α is of order 2, the map $\alpha\colon S^4\to Y$ extends to a map $\tilde{\alpha}\colon M(\mathbb{Z}/2,4)\to Y$ with

$$\tilde{\alpha}_*\colon H_4(M(\mathbb{Z}/2,4))\longrightarrow H_4(Y)$$

an isomorphism. Let

$$j\colon M(\mathbb{Z}/2,3)\longrightarrow Y$$

be the canonical inclusion. Then $j_*\colon H_3(M(\mathbb{Z}/2,3))\to H_3(Y)$ is an isomorphism. Then

$$(j,\tilde{\alpha})\colon M(\mathbb{Z}/2,3)\vee M(\mathbb{Z}/2,4)\longrightarrow Y$$

is a homotopy equivalence because it induces an isomorphism on homology, which contradicts that the Steenrod operation $Sq^2\colon H^3(Y;\mathbb{Z}/2)\to H^5(Y;\mathbb{Z}/2)$ is an isomorphism. Thus $\pi_4(Y)=\mathbb{Z}/4$.

Now the assertion follows from decomposition (3.6) and the computational formulae (3.8) and (3.9). □

Corollary 3.2. *Let A_2 be any elementary 2-group. Then there is a natural short exact sequence*

$$(A_2 \otimes \Lambda^2(A_2))^{\oplus 2} \hookrightarrow \pi_4(\Sigma K(A_2,1)) \twoheadrightarrow \frac{1}{2}(A_2 \otimes A_2),$$

where $\frac{1}{2}(A_2 \otimes A_2)$ is a free $\mathbb{Z}/4$-module. Moreover this splits off unnaturally.

Proof. By Theorem 3.1, there is a natural short exact sequence

$$(A_2 \otimes \Lambda^2(A_2))^{\oplus 2} \hookrightarrow \pi_4(\Sigma K(A_2,1)) \twoheadrightarrow \pi_4(\Sigma K(A_2,1))/(A_2 \otimes \Lambda^2(A_2))^{\oplus 2}.$$

By Theorem 3.2, the quotient group $\pi_4(\Sigma K(A_2,1))/(A_2 \otimes \Lambda^2(A_2))^{\oplus 2}$ is free $\mathbb{Z}/4$-module for any finite dimensional elementary 2-groups. The assertion follows by taking direct limits. □

Remark 3.1. *The summand $\frac{1}{2}(A_2 \otimes A_2)$ is sub-quotient functor of $\pi_4(\Sigma K(A,1))$ on A in the following sense. For any abelian group A, the $\mathbb{Z}/2$-component A_2 is given by the image of*

$$Sq^1_*\colon H_2(A;\mathbb{Z}/2) \longrightarrow H_1(A;\mathbb{Z}/2).$$

Thus $A \mapsto A_2$ is a sub functor of the identity functor on abelian groups. Then $\pi_4(\Sigma K(A_2,1))$ is a sub functor of $\pi_4(\Sigma K(A,1))$ on A and so

$$\frac{1}{2}(A_2 \otimes A_2) = \pi_4(\Sigma K(A_2,1))/(A_2 \otimes \Lambda^2(A_2))^{\oplus 2}$$

is a sub-quotient functor of $\pi_4(\Sigma K(A,1))$ on A.

3.2. **Applications.** As an application, we compute $\pi_i(M(\mathbb{Z}/p^r,2))$ for $i \leq 4$. By the Hurewicz Theorem, $\pi_2(M(\mathbb{Z}/p^r,2)) = \mathbb{Z}/p^r$. From the Whitehead exact sequence (2.2), we have

$$\Gamma_n(M(\mathbb{Z}/p^r,2)) = \pi_n(M(\mathbb{Z}/p^r,2)) \tag{3.10}$$

for $r \geq 3$ because $H_i(M(\mathbb{Z}/p^r,2) = 0$ for $i \geq 3$. It follows directly that

$$\pi_3(M(\mathbb{Z}/p^r,2)) = \Gamma_3(M(\mathbb{Z}/p^r,2)) = \Gamma_2(\mathbb{Z}/p^r) = \begin{cases} \mathbb{Z}/p^r & \text{if} \quad p > 2, \\ \mathbb{Z}/2^{r+1} & \text{if} \quad p = 2, \end{cases} \tag{3.11}$$

where $\Gamma_2(A)$ is computed in Example 2.1. From Theorem 2.2 (1), there is a short exact sequence

$$\Gamma_2^2(M(\mathbb{Z}/p^r,2)) \hookrightarrow \Gamma_4(M(\mathbb{Z}/p^r,2)) \twoheadrightarrow R_2(\pi_2(M(\mathbb{Z}/p^r,2))) = R_2(\mathbb{Z}/p^r).$$

From Subsection 2.2,

$$R_2(\mathbb{Z}/p^r) = \begin{cases} 0 & \text{if} \quad p > 2, \\ \mathbb{Z}/2 & \text{if} \quad p = 2. \end{cases}$$

By the definition (2.4) of the functor Γ_2^2, the group $\Gamma_2^2(M(\mathbb{Z}/p^r,2))$ is given by the push-out

$$\begin{tikzcd}
\Gamma_2(\mathbb{Z}/p^r) \otimes (\mathbb{Z}/p^r \oplus \mathbb{Z}/2) \arrow[r, "q\oplus \mathrm{id}"] \arrow[d, "\eta_1\otimes\mathrm{id}", "\cong"'] & \mathcal{L}_s^3(\mathbb{Z}/p^r) \oplus \Gamma_2(\mathbb{Z}/p^r) \otimes \mathbb{Z}/2 \arrow[d] \\
\pi_3(M(\mathbb{Z}/p^r,2)) \otimes (\mathbb{Z}/p^r \oplus \mathbb{Z}/2) \arrow[r] & \Gamma_2^2(M(\mathbb{Z}/p^r,2)).
\end{tikzcd}$$

Thus

$$\Gamma_2^2(M(\mathbb{Z}/p^r,2)) \cong \mathcal{L}_s^3(\mathbb{Z}/p^r) \oplus \Gamma_2(\mathbb{Z}/p^r) \otimes \mathbb{Z}/2.$$

Since $\mathbb{Z}/p^r$ is cyclic and $\mathcal{L}_s^3(A)$ is isomorphic to the kernel of $A\otimes\Lambda^2(A)\to\Lambda^3(A)$, we have

$$\mathcal{L}_s^3(\mathbb{Z}/p^r)=0$$

and so

$$\Gamma_2^2(M(\mathbb{Z}/p^r,2))=\begin{cases}0 & \text{if} \quad p>2,\\ \mathbb{Z}/2 & \text{if} \quad p=2.\end{cases}$$

A direct consequence is:

$$\pi_4(\Sigma M(\mathbb{Z}/p^r,2))=0 \text{ for } p>2. \tag{3.12}$$

For the case $p=2$, we have the short exact sequence

$$\mathbb{Z}/2 \hookrightarrow \pi_4(M(\mathbb{Z}/2^r,2)) \twoheadrightarrow \mathbb{Z}/2.$$

The remaining problem is to decide whether $\pi_4(M(\mathbb{Z}/2^r,2))$ is equal to $\mathbb{Z}/2\oplus\mathbb{Z}/2$ or $\mathbb{Z}/4$. It has been computed in [22] that $\pi_4(M(\mathbb{Z}/2,2))=\mathbb{Z}/4$. For $r>1$, the group $\pi_4(M(\mathbb{Z}/2^r,2))$ seems not recorded in references. We are going to determine the group $\pi_4(M(\mathbb{Z}/2^r,2))$ using our methods.

Lemma 3.5. *Let*

$$j\colon M(\mathbb{Z}/2^r,2)\longrightarrow \Sigma K(\mathbb{Z}/2^r,1)$$

be the canonical map inducing isomorphism on H_2. *Then*

$$j_*\colon \Gamma_4(M(\mathbb{Z}/2^r,2))\longrightarrow \Gamma_4(\Sigma K(\mathbb{Z}/2^r,1))$$

is an isomorphism.

Proof. By Theorem 2.2(1), there is a commutative diagram of short exact sequences

$$\begin{tikzcd}
\Gamma_2^2(M(\mathbb{Z}/2^r,2))=\mathbb{Z}/2 \arrow[r, hook] \arrow[d, "j_*"] & \Gamma_4(M(\mathbb{Z}/2^r,2)) \arrow[r, two heads] \arrow[d, "j_*"] & R_2(\mathbb{Z}/2^r)=\mathbb{Z}/2 \arrow[d, "j_*", "\cong"'] \\
\Gamma_2^2(K(\mathbb{Z}/2^r,2)) \arrow[r, hook] & \Gamma_4(K(\mathbb{Z}/2^r,2)) \arrow[r, two heads] & R_2(\mathbb{Z}/2^r)=\mathbb{Z}/2
\end{tikzcd}$$

From the Whitehead exact sequence

$$\Gamma_3(\mathbb{Z}/2^r)=\mathbb{Z}/2^{r+1}\to\pi_3(\Sigma K(\mathbb{Z}/2^r,1))=\mathbb{Z}/2^r\otimes\mathbb{Z}/2^r=\mathbb{Z}/2^r\to 0,$$

we have

$$\eta_1\otimes\mathrm{id}\colon \Gamma_2(\mathbb{Z}/2^r)\otimes(\mathbb{Z}/2^r\oplus\mathbb{Z}/2)\longrightarrow \pi_3(K(\mathbb{Z}/2^r,2))\otimes(\mathbb{Z}/2^r\oplus\mathbb{Z}/2)$$

is an isomorphism. Similar to the computation of $\Gamma_2^2(M(\mathbb{Z}/2^r,2))$, we have

$$\Gamma_2^2(K(\mathbb{Z}/2^r,2))=\mathbb{Z}/2$$

with an isomorphism $j_*\colon \Gamma_2^2(M(\mathbb{Z}/2^r,2))\cong\Gamma_2^2(K(\mathbb{Z}/2^r,2))$. The assertion then follows by 5-lemma. □

Lemma 3.6. *The group*

$$\Gamma_4(\Sigma K(\mathbb{Z}/2^r,1))=\begin{cases}\mathbb{Z}/2\oplus\mathbb{Z}/2 & \textit{if} \quad r>1,\\ \mathbb{Z}/4 & \textit{if} \quad r=1.\end{cases}$$

Proof. Let $Z=\Sigma K(\mathbb{Z}/2^r,1)\wedge K(\mathbb{Z}/2^r,1)$ and let $H\colon Z\to\Sigma K(\mathbb{Z}/2^r,1)$ be the Hopf map. From equation (3.5), there is a commutative diagram of exact sequence

$$\begin{tikzcd}
\Gamma_4(Z)=\mathbb{Z}/2 \arrow[r, hook] \arrow[d, "H_*"] & \pi_4(Z) \arrow[r, two heads] \arrow[d, "H_*", "\cong"'] & H_4(Z)=\mathbb{Z}/2^r \arrow[d, "H_*"] \\
\Gamma_4(\Sigma K(\mathbb{Z}/2^r,1)) \arrow[r, hook] & \pi_4(\Sigma K(\mathbb{Z}/2^r,1)) \arrow[r] & H_4(\Sigma K(\mathbb{Z}/2^r,1)),
\end{tikzcd}$$

where the bottom row is left exact because $H_5(\Sigma K(\mathbb{Z}/2^r,1))=H_4(\mathbb{Z}/2^r)=0$.

If $r>1$, then $\Gamma_4(Z)$ is a summand of $\pi_4(Z)\cong\pi_4(\Sigma K(\mathbb{Z}/2^r,1))$ by Theorem 3.2. Thus $\Gamma_4(Z)=\mathbb{Z}/2$ is also a summand of $\Gamma_4(\Sigma K(\mathbb{Z}/2^r,1))$. It follows that

$$\Gamma_4(\Sigma K(\mathbb{Z}/2^r,1))=\mathbb{Z}/2\oplus\mathbb{Z}/2 \text{ if } r>1.$$

If $r=1$, by Corollary 3.2, $\pi_4(\Sigma K(\mathbb{Z}/2,1))=\mathbb{Z}/4$ and so

$$\Gamma_4(\Sigma K(\mathbb{Z}/2,1))\cong\pi_4(\Sigma K(\mathbb{Z}/2^r,1))=\mathbb{Z}/4.$$

The proof is finished. □

Since $\Gamma_4(\Sigma K(\mathbb{Z}/2^r,1))\to\pi_4(\Sigma K(\mathbb{Z}/2^r,1))$ is a monomorphism, from Lemmas 3.5 and 3.6, we have the following:

Corollary 3.3. *Let*

$$j\colon M(\mathbb{Z}/2^r,2)\longrightarrow\Sigma K(\mathbb{Z}/2^r,1)$$

be the canonical map inducing isomorphism on H_2*. Then*

1. $\pi_4(M(\mathbb{Z}/2,2))=\mathbb{Z}/4$ *and*

   $$j_*\colon\pi_4(M(\mathbb{Z}/2,2))\to\pi_4(\Sigma K(\mathbb{Z}/2,1))$$

   is an isomorphism.
2. *For* $r>1$, $\pi_4(M(\mathbb{Z}/2^r,2))=\mathbb{Z}/2\oplus\mathbb{Z}/2$ *and*

   $$j_*\colon\pi_4(M(\mathbb{Z}/2^r,2))=\mathbb{Z}/2\oplus\mathbb{Z}/2\to\pi_4(\Sigma K(\mathbb{Z}/2^r,1))=\mathbb{Z}/2\oplus\mathbb{Z}/2^r$$

   is a monomorphism. □

Note that $M(\mathbb{Z}/2,2)=\Sigma\mathbb{R}\mathrm{P}^2$ and $\Sigma K(\mathbb{Z}/2,1)=\Sigma\mathbb{R}\mathrm{P}^\infty$ with the canonical inclusion $j\colon\Sigma\mathbb{R}\mathrm{P}^2\hookrightarrow\Sigma\mathbb{R}\mathrm{P}^\infty$. A consequence of Corollary 3.3 (1) on the suspended projective spaces are as follows.

Corollary 3.4. *Let* $j\colon\Sigma\mathbb{R}\mathrm{P}^2\to\Sigma\mathbb{R}\mathrm{P}^n$ *be the canonical inclusion with* $3\leq n\leq\infty$*.*

1. *For* $4\leq n\leq\infty$, $j_*\colon\pi_4(\Sigma\mathbb{R}\mathrm{P}^2)=\mathbb{Z}/4\to\pi_4(\Sigma\mathbb{R}\mathrm{P}^n)$ *is an isomorphism.*
2. *For* $n=3$, $j_*\colon\pi_4(\Sigma\mathbb{R}\mathrm{P}^2)=\mathbb{Z}/4\to\pi_4(\Sigma\mathbb{R}\mathrm{P}^3)$ *is a splitting monomorphism. Moreover*

   $$\pi_4(\Sigma\mathbb{R}\mathrm{P}^3)\cong\pi_4(\Sigma\mathbb{R}\mathrm{P}^2)\oplus\mathbb{Z}=\mathbb{Z}/4\oplus\mathbb{Z}.$$

Proof. Assertion (1) and the first part of assertion (2) are direct consequences of Corollary 3.3. For the second part of assertion (2), notice that $\mathbb{R}\mathrm{P}^3=SO(3)$. From the commutative diagram

$$\begin{tikzcd}
\pi_4(\Sigma SO(3)\wedge SO(3)) \arrow[r, "\cong"] \arrow[d, "H_*"] & \pi_4(\Sigma\mathbb{R}\mathrm{P}^\infty\wedge\mathbb{R}\mathrm{P}^\infty) \arrow[d, "H_*", "\cong"'] \\
\pi_4(\Sigma SO(3)) \arrow[r] & \pi_4(\Sigma\mathbb{R}\mathrm{P}^\infty),
\end{tikzcd}$$

we have

$$\begin{aligned}
\pi_4(\Sigma SO(3)) &\cong \pi_4(\Sigma SO(3)\wedge SO(3))\oplus\pi_4(BSO(3))\\
&\cong \pi_4(\Sigma\mathbb{R}\mathrm{P}^\infty)\oplus\pi_3(SO(3))\\
&\cong \mathbb{Z}/4\oplus\mathbb{Z}
\end{aligned}$$

and hence the result. □

Another consequence is as follows:

Corollary 3.5. *Let Σ_3 be the third symmetric group. Then $\pi_4(\Sigma K(\Sigma_3,1))=\mathbb{Z}/12$.*

Proof. Recall that the integral homology groups of Σ_3 are 4-periodic with the following initial terms:

$$H_1(\Sigma_3)=\mathbb{Z}/2,\ H_2(\Sigma_3)=0,\ H_3(\Sigma_3)=\mathbb{Z}/6,\ H_4(\Sigma_3)=0.$$

Let $X=\Sigma K(\Sigma_3,1)$. The Whitehead exact sequence has the following form:

$$\begin{tikzcd}
\Gamma_4(X) \arrow[r, hook] & \pi_4(X) \arrow[r] & H_3(\Sigma_3)=\mathbb{Z}/6 \arrow[r] & \Gamma_3(X)=\mathbb{Z}/4 \arrow[r, two heads] & \pi_3(X)=\mathbb{Z}/2.
\end{tikzcd}$$

The inclusion $\Sigma_2=\mathbb{Z}/2\to\Sigma_3$ induces an isomorphism

$$\pi_i(\Sigma K(\mathbb{Z}/2,1))=\mathbb{Z}/2\xrightarrow{\cong}\pi_i(\Sigma K(\Sigma_3,1))=\mathbb{Z}/2$$

for $i=2,3$. By Theorem 2.2 (1) together with Lemma 3.6, the inclusion $\Sigma_2=\mathbb{Z}/2\to\Sigma_3$ induces an

$$\Gamma_4(\Sigma K(\mathbb{Z}/2,1))=\mathbb{Z}/4\xrightarrow{\cong}\Gamma_4(\Sigma K(\Sigma_3,1))$$

and hence the result. □

4. On group $\pi_5(\Sigma K(A,1))$

4.1. Some Properties of the Functor $A\mapsto\pi_5(\Sigma K(A,1))$.

From Hopf fibration

$$\Sigma K(A,1)\wedge K(A,1)\longrightarrow\Sigma K(A,1)\longrightarrow K(A,2),$$

it suffices to compute $\pi_5(\Sigma K(A,1)\wedge K(A,1))$. Let $Z=\Sigma K(A,1)\wedge K(A,1)$. Since Z is 2-connected, from Theorem 2.2(2), there are natural exact sequences

$$\begin{tikzcd}
 & \pi_4(\Sigma K(A,1))\otimes\mathbb{Z}/2\oplus\Lambda^2(A\otimes A) \arrow[d, hook] & & \\
H_6(Z) \arrow[r] & \Gamma_5(Z) \arrow[r] \arrow[d, two heads] & \pi_5(Z) \arrow[r, two heads] & H_5(Z), \\
 & \mathrm{Tor}(A\otimes A,\mathbb{Z}/2) & &
\end{tikzcd} \tag{4.1}$$

where $\pi_5(Z)\to H_5(Z)$ is onto by equation (3.5). The group $\pi_4(\Sigma K(A,1))$ has been determined by Theorems 3.1 and 3.2.

Proposition 4.1. *Let A be a free abelian group. Then there is a natural short exact sequence*

$$(\Lambda^2(A)\otimes A\otimes\mathbb{Z}/2)^{\oplus 2}\oplus A^{\otimes 2}\otimes\mathbb{Z}/2\oplus\Lambda^2(A\otimes A)\hookrightarrow\pi_5(\Sigma K(A,1))\twoheadrightarrow(\Lambda^3(A)\otimes A)^{\oplus 2}\oplus\Lambda^2(A)^{\otimes 2}.$$

Proof. Since A is a free abelian group, the Hurewicz homomorphism $h_*\colon\pi_*(Z)\to\tilde{H}_*(Z)$ is onto because Z is a wedge of spheres. Thus there is a short exact sequence

$$\Gamma_5(Z)\hookrightarrow\pi_5(Z)\twoheadrightarrow H_5(Z).$$

By Theorem 3.1, $\pi_4(Z)\cong(\Lambda^2(A)\otimes A)^{\oplus 2}\oplus A\otimes A\otimes\mathbb{Z}/2$ for a free abelian group A. The assertion follows from diagram (4.1). □

Proposition 4.2. *If A is a torsion abelian group with the property that $2\colon A\to A$ is an isomorphism, then there is a natural short exact sequence*

$$\Lambda^2(A\otimes A)\hookrightarrow\pi_5(\Sigma K(A,1))\twoheadrightarrow H_4(K(A,1)\wedge K(A,1)).$$

Proof. It suffices to show that the Hurewicz homomorphism $h_*\colon\pi_6(Z)\to H_6(Z)$ is onto. We may assume that A is finitely generated because we can take direct limit for general case whence the finitely generated case is proved. Then A is a direct sum of the primary p-torsion groups $\mathbb{Z}/p^r$ for some $r\geq 1$ and odd primes p. According to [12], there is homotopy decomposition

$$\Sigma K(\mathbb{Z}/p^r,1)\simeq X_1\vee\cdots\vee X_{p-1},$$

where $\bar{H}_q(X_i;\mathbb{Z})\neq 0$ if and only if $q\equiv 2i \mod 2p-2$. Together with the decomposition formula (3.7) for the smash product of Moore spaces, up to 6-skeleton, $\Sigma K(A,1)\wedge K(A,1)$ is homotopy equivalent to a wedge of spheres and Moore spaces. It follows that the Hurewicz homomorphism

$$\pi_6(\Sigma K(A,1)\wedge K(A,1))\longrightarrow H_6(\Sigma K(A,1)\wedge K(A,1))$$

is onto and hence the result. □

From the above proof, we also have the following:

Proposition 4.3. *Let A be any abelian group. Let $\mathbb{Z}_{\frac{1}{2}}=\{\frac{m}{2^r}\in\mathbb{Q}\mid m\in\mathbb{Z},\ r\geq 0\}$. Then there is natural short exact sequence*

$$\Lambda^2(A\otimes A)\otimes\mathbb{Z}_{\frac{1}{2}}\hookrightarrow\pi_5(\Sigma K(A,1))\otimes\mathbb{Z}_{\frac{1}{2}}\twoheadrightarrow H_4(K(A,1)\wedge K(A,1))\otimes\mathbb{Z}_{\frac{1}{2}}.$$

□

For computing the group $\pi_5(\Sigma K(A,1))$, as one see from the above, the tricky part is the 2-torsion. Whence A contains 2-torsion summands, the Hurewicz homomorphism $\pi_6(Z)\to H_6(Z)$ is no longer epimorphism in general and so $\Gamma_5(Z)\to\pi_5(Z)$ is not a monomorphism in general. Also the group $\pi_5(Z)$ in diagram (4.1) admits non-trivial extension. The computation of the group $\pi_5(\Sigma K(A,1))$ for finitely generated abelian groups A can be given by the following steps:

Step 1. Take a primary decomposition of A and write $K(A,1)$ as a product of copies of $S^1=K(\mathbb{Z},1)$ and $K(\mathbb{Z}/p^r,1)$.

Step 2. By using the fact that $\Sigma X\times Y\simeq \Sigma X\vee\Sigma Y\simeq\Sigma X\wedge Y$ for any spaces X and Y, one gets

$$\begin{aligned}\Sigma(X_1\times X_2)\wedge(X_1\times X_2) &\simeq \Sigma(X_1\vee X_2\vee X_1\wedge X_2)\wedge(X_1\vee X_2\vee X_1\wedge X_2)\\ &\simeq \Sigma(X_1^{\wedge 2}\vee X_2^{\wedge 2}\vee X_1^{\wedge 2}\wedge X_2^{\wedge 2}\vee\\ &\qquad \bigvee^{2} X_1\wedge X_2\vee\bigvee^{2} X_1^{\wedge 2}\wedge X_2\vee\bigvee^{2} X_1\wedge X_2^{\wedge 2}).\end{aligned}$$

From this, $\Sigma K(A,1)\wedge K(A,1)$ is then homotopy equivalent to a wedge of the spaces in the form

$$X=\Sigma^m K(\mathbb{Z}/p_1^{r_1},1)\wedge K(\mathbb{Z}/p_2^{r_2},1)\wedge\cdots\wedge K(\mathbb{Z}/p_t^{r_t},1)$$

with $m+t\geq 3$ and $m\geq 1$.

Step 3. By applying the Hilton-Milnor Theorem, we have

$$\begin{aligned}\Omega(\Sigma X\vee\Sigma Y) &\simeq \Omega\Sigma X\times\Omega\Sigma Y\times\Omega\Sigma((\Omega\Sigma X)\wedge(\Omega\Sigma Y))\\ &\simeq \Omega\Sigma X\times\Omega\Sigma Y\times\Omega\Sigma\left(\bigvee_{i,j=1}^{\infty}X^{\wedge i}\wedge Y^{\wedge j}\right).\end{aligned}$$

Thus

$$\pi_n(\Sigma X\vee\Sigma Y)\cong\pi_n(\Sigma X)\oplus\pi_n(\Sigma Y)\oplus\pi_n\left(\bigvee_{i,j=1}^{\infty}\Sigma X^{\wedge i}\wedge Y^{\wedge j}\right).$$

Note that the connectivity of $X^{\wedge i}\wedge Y^{\wedge j}$ tends to ∞ as $i,j\to\infty$. By repeating the above procedure, $\pi_n(\Sigma K(A,1)\wedge K(A,1))$ is isomorphism to a direct sum of the groups $\pi_n(X)$ with X given in the form above.

Notice that

$$\Sigma^m K(\mathbb{Z}/p_1^{r_1},1)\wedge K(\mathbb{Z}/p_2^{r_2},1)\wedge\cdots\wedge K(\mathbb{Z}/p_t^{r_t},1)\simeq *$$

if the primes $p_i\neq p_j$ for some $i\neq j$. Thus we only need to compute

$$\pi_5(\Sigma^m K(\mathbb{Z}/p^{r_1},1)\wedge K(\mathbb{Z}/p^{r_2},1)\wedge\cdots\wedge K(\mathbb{Z}/p^{r_t},1))$$

for a prime p. If $t=0$, the homotopy group $\pi_5(S^m)$ is known by $\pi_5(S^3)=\pi_5(S^4)=\mathbb{Z}/2$ and $\pi_5(S^5)=\mathbb{Z}$. For an odd prime p, this homotopy group can be determined by Proposition 4.2. The rest work in this section is of course to compute $\pi_5(\Sigma^m K(\mathbb{Z}/2^{r_1},1)\wedge K(\mathbb{Z}/2^{r_2},1)\wedge\cdots\wedge K(\mathbb{Z}/2^{r_t},1))$ with $m+t\geq 3$. When $m+t\geq 5$, we have

$$\pi_5(X)=\begin{cases}0 & \text{if}\quad m+t>5\\ \mathbb{Z}/2^{\min\{r_1,\ldots,r_t\}} & \text{if}\quad m+t=5\text{ with }t\geq 1\end{cases}$$

for $X=\Sigma^m K(\mathbb{Z}/2^{r_1},1)\wedge K(\mathbb{Z}/2^{r_2},1)\wedge\cdots\wedge K(\mathbb{Z}/2^{r_t},1)$. The first less obvious case is $m+t=4$, which will be discussed in the next subsection.

4.2. **The Group $\pi_5(\Sigma^m K(\mathbb{Z}/2^{r_1},1)\wedge K(\mathbb{Z}/2^{r_2},1)\wedge\cdots\wedge K(\mathbb{Z}/2^{r_t},1))$ for $m+t=4$ and $m,t\geq 1$.**

We first consider the case $t=1$.

Lemma 4.1. *The Hurewicz homomorphism*

$$h_5\colon\pi_5(\Sigma^2 K(\mathbb{Z}/2^r,1))\to H_5(\Sigma^2 K(\mathbb{Z}/2^r,1))$$

is onto for any $r\geq 1$.

Proof. Let $X=\Sigma^2K(\mathbb{Z}/2^r,1)$. Consider the Whitehead exact sequence

$$\pi_5(X)\xrightarrow{h_5}H_5(X)\to\Gamma_4(X)=\mathbb{Z}/2\to\pi_4(X)\to H_4(X)=0.$$

Thus the Hurewicz homomorphism h_5 is onto if and only if $\pi_4(X)\neq 0$.

Let $f\colon S^3\to X$ be a map representing the generator for $\pi_3(X)=\mathbb{Z}/2^r$. From the remark to Theorem 2.1, $\pi_4(X)=0$ if and only if the composite

$$S^4\xrightarrow{\eta}S^3\xrightarrow{f}X$$

is null homotopic, if and only if the map $f\colon S^3\to X$ extends to a map $\tilde{f}\colon\Sigma\mathbb{C}\mathrm{P}^2\to X$ because $\Sigma\mathbb{C}\mathrm{P}^2$ is the homotopy cofibre of $\eta\colon S^4\to S^3$.

Suppose that there exists a map $\tilde{f}\colon\Sigma\mathbb{C}\mathrm{P}^2\to X$ such that $\tilde{f}|_{S^4}=f$. By taking mod 2 cohomology, there is commutative diagram

$$\begin{tikzcd}
H^5(\Sigma\mathbb{C}\mathrm{P}^2;\mathbb{Z}/2) & H^5(X;\mathbb{Z}/2)=\mathbb{Z}/2 \arrow[l, "\tilde{f}^*"'] \\
H^3(\Sigma\mathbb{C}\mathrm{P}^2;\mathbb{Z}/2)=H^3(S^3;\mathbb{Z}/2) \arrow[u, "\cong", "Sq^2"'] & H^3(X;\mathbb{Z}/2)=\mathbb{Z}/2. \arrow[l, "\tilde{f}^*=f^*"', "\cong"] \arrow[u, "Sq^2"']
\end{tikzcd}$$

It follows that

$$Sq^2\colon H^3(X;\mathbb{Z}/2)\longrightarrow H^5(X;\mathbb{Z}/2)$$

is an isomorphism. On the other hand, from the fact that $X=\Sigma^2K(\mathbb{Z}/2^r,1)$ and $Sq^2\colon H^1(K(\mathbb{Z}/2^r,1);\mathbb{Z}/2)\to H^3(K(\mathbb{Z}/2^r,1))$ is zero, $Sq^2\colon H^3(X;\mathbb{Z}/2)\to H^5(X;\mathbb{Z}/2)$ is zero. This gives a contradiction. The assertion follows. ☐

Proposition 4.4. $\pi_5(\Sigma^3K(\mathbb{Z}/2^r,1))=\mathbb{Z}/2$ *for* $r\geq 1$.

Proof. Let $X=\Sigma^3K(\mathbb{Z}/2^r,1)$. Consider the Whitehead exact sequence

$$\pi_6(X)\xrightarrow{h_6}H_6(X)\to\Gamma_5(X)=\mathbb{Z}/2\to\pi_5(X)\to H_5(X)=0.$$

By Lemma 4.1, $h_6\colon\pi_6(X)\to H_6(X)$ is onto. Thus $\pi_5(X)\cong\Gamma_5(X)=\mathbb{Z}/2$. ☐

Now we consider the case $t=2$.

Proposition 4.5. *Let* $r_1,r_2\geq 1$. *Then*

$$\pi_5(\Sigma^2K(\mathbb{Z}/2^{r_1},1)\wedge K(\mathbb{Z}/2^{r_2},1))=\begin{cases}\mathbb{Z}/2\oplus\mathbb{Z}/2^{\min\{r_1,r_2\}} & \text{if}\quad \max\{r_1,r_2\}>1,\\ \mathbb{Z}/4 & \text{if}\quad r_1=r_2=1.\end{cases}$$

Proof. Let $X=\Sigma^2K(\mathbb{Z}/2^{r_1},1)\wedge K(\mathbb{Z}/2^{r_2},1)$. By Lemma 4.1, there exists a map

$$f_i\colon S^5\longrightarrow\Sigma^2K(\mathbb{Z}/2^{r_i},1),\quad i=1,2,$$

which induces an epimorphism

$$f_{i*}\colon H_5(S^5)\twoheadrightarrow H_5(\Sigma^2K(\mathbb{Z}/2^{r_i},1).$$

Let $j\colon Y=\Sigma^2M(\mathbb{Z}/2^{r_1},1)\wedge M(\mathbb{Z}/2^{r_2},1)\hookrightarrow X$ be the canonical inclusion. Then the map

$$f\colon Y\vee S^5\wedge K(\mathbb{Z}/2^{r_2},1)\vee K(\mathbb{Z}/2^{r_1},1)\wedge S^5\xrightarrow{(j,f_1\wedge\mathrm{id},\mathrm{id}\wedge f_2)}X$$

induces an isomorphism on $H_j(\ ;\mathbb{Z}/2)$ for $j\leq 6$. Thus

$$f_*\colon\pi_k\left(Y\vee S^5\wedge K(\mathbb{Z}/2^{r_2},1)\vee K(\mathbb{Z}/2^{r_1},1)\wedge S^5\right)\longrightarrow\pi_k(X)$$

is an isomorphism for $k \leq 5$. Note that

$$\pi_k(S^5 \wedge K(\mathbb{Z}/2^{r_2}, 1)) = \pi_k(K(\mathbb{Z}/2^{r_1}, 1) \wedge S^5) = 0$$

for $k \leq 5$. Thus

$$j_*\colon \pi_k(Y) \to \pi_k(X)$$

is an isomorphism for $k \leq 5$. In particular, $\pi_5(Y) \cong \pi_5(X)$.

If $\max r_1, r_2 > 1$, from decomposition (3.7), we have

$$\Sigma^2 M(\mathbb{Z}/2^{r_1}, 1) \wedge M(\mathbb{Z}/2^{r_2}, 1) \simeq M(\mathbb{Z}/2^{\min\{r_1,r_2\}}, 4) \vee M(\mathbb{Z}/2^{\min\{r_1,r_2\}}, 5)$$

and so

$$\pi_5(Y) \cong \pi_5(M(\mathbb{Z}/2^{\min\{r_1,r_2\}}, 4)) \oplus \pi_5(M(\mathbb{Z}/2^{\min\{r_1,r_2\}}, 5)) = \mathbb{Z}/2 \oplus \mathbb{Z}/2^{\min\{r_1,r_2\}}.$$

Consider the case $r_1 = r_2 = 1$. From formula (3.9) and the Freudenthal Suspension Theorem,

$$\pi_5(\Sigma^2 M(\mathbb{Z}/2, 1) \wedge M(\mathbb{Z}/2, 1)) \cong \pi_4(\Sigma M(\mathbb{Z}/2, 1) \wedge M(\mathbb{Z}/2, 1)) \cong \mathbb{Z}/4.$$

The proof is finished. □

The last case is $t = 3$.

Proposition 4.6. *Let $r_1, r_2, r_3 \geq 1$ and let $r = \min\{r_1, r_2, r_3\}$. Then*

$$\pi_5(\Sigma K(\mathbb{Z}/2^{r_1}, 1) \wedge K(\mathbb{Z}/2^{r_2}, 1) \wedge K(\mathbb{Z}/2^{r_3}, 1)) = \begin{cases} \mathbb{Z}/2 \oplus \mathbb{Z}/2^r & \text{if} \quad \max\{r_1, r_2, r_3\} > 1, \\ \mathbb{Z}/2 \oplus \mathbb{Z}/2 & \text{if} \quad r_1 = r_2 = r_3 = 1. \end{cases}$$

Proof. Let $X = \Sigma K(\mathbb{Z}/2^{r_1}, 1) \wedge K(\mathbb{Z}/2^{r_2}, 1) \wedge K(\mathbb{Z}/2^{r_3}, 1)$. Let f_1 be the composite

$$S^6 \xrightarrow{g} \Sigma^3 K(\mathbb{Z}/2^{r_1}, 1) \cong \Sigma K(\mathbb{Z}/2^{r_1}, 1) \wedge S^1 \wedge S^1 \hookrightarrow X,$$

where g is a map which induces epimorphism on $H_6(\ )$ by Lemma 4.1. Similarly, we have the maps

$$f_i\colon S^6 \longrightarrow \Sigma K(\mathbb{Z}/2^{r_1}, 1) \wedge K(\mathbb{Z}/2^{r_2}, 1) \wedge K(\mathbb{Z}/2^{r_3}, 1), \quad i = 2, 3,$$

by replacing $K(\mathbb{Z}/2^{r_1}, 1)$ by $K(\mathbb{Z}/2^{r_i}, 1)$. Let $Y = \Sigma M(\mathbb{Z}/2^{r_1}, 1) \wedge M(\mathbb{Z}/2^{r_2}, 1) \wedge M(\mathbb{Z}/2^{r_3}, 1)$ and let $j\colon Y \hookrightarrow X$ be the canonical inclusion. The map

$$Y \vee S^6 \vee S^6 \vee S^6 \xrightarrow{(j, f_1, f_2, f_3)} X$$

induces an isomorphism on $H_k(\ ; \mathbb{Z}/2)$ for $k \leq 6$ and so

$$(j, f_1, f_2, f_3)_*\colon \pi_5(Y \vee S^6 \vee S^6 \vee S^6) = \pi_5(Y) \longrightarrow \pi_5(X)$$

is an isomorphism.

If $\max\{r_1, r_2, r_3\} > 1$, from decomposition (3.7),

$$Y \simeq M(\mathbb{Z}/2^r, 4) \vee M(\mathbb{Z}/2^r, 5) \vee M(\mathbb{Z}/2^r, 5) \vee M(\mathbb{Z}/2^r, 6)$$

and so

$$\pi_5(Y) = \mathbb{Z}/2 \oplus \mathbb{Z}/2^r \oplus \mathbb{Z}/2^r.$$

If $r_1 = r_2 = r_3 = 1$, there is a homotopy decomposition [22, Corollary 3.7]

$$\Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2 \simeq \Sigma\mathbb{C}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2 \vee \Sigma^4\mathbb{R}\mathrm{P}^2 \vee \Sigma^4\mathbb{R}\mathrm{P}^2.$$

By [22, Lemma 6.34 (2)],

$$\pi_5(\Sigma\mathbb{C}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2) = 0$$

and so

$$\pi_5(\Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2) = \mathbb{Z}/2 \oplus \mathbb{Z}/2,$$

which finishes the proof. □

Remark 4.1. For the case $X = \Sigma K(\mathbb{Z}/2,1) \wedge K(\mathbb{Z}/2,1) \wedge K(\mathbb{Z}/2,1)$, the Hurewicz homomorphism

$$\pi_5(X) = \mathbb{Z}/2 \oplus \mathbb{Z}/2 \longrightarrow H_5(X) = \mathbb{Z}/2 \oplus \mathbb{Z}/2$$

is an isomorphism and so, in the Whitehead exact sequence,

$$H_6(X) \longrightarrow \Gamma_5(X) = \mathbb{Z}/2$$

is onto. This gives an example that the morphism $H_6(Z) \to \Gamma_5(Z)$ in diagram (4.1) may not be zero, which is the only example in the case $m+t=4$. More examples will be shown up in the case $m+t=3$ in the next subsections.

4.3. **The Group** $\pi_5(\Sigma K(\mathbb{Z}/2^r,1)) \cong \pi_5(\Sigma K(\mathbb{Z}/2^r,1) \wedge K(\mathbb{Z}/2^r,1))$.

Lemma 4.2. *Let* $X = \Sigma K(\mathbb{Z}/2^r,1) \wedge K(\mathbb{Z}/2^r,1)$ *with* $r \geq 1$. *Then mod 2 Hurewicz homomorphism*

$$\pi_6(X) \xrightarrow{h_6} H_6(X) \longrightarrow H_6(X;\mathbb{Z}/2)$$

is zero.

Proof. Recall that the mod 2 cohomology ring

$$H^*(K(\mathbb{Z}/2^r,1);\mathbb{Z}/2) \cong E(u_1) \otimes P(u_2)$$

with the rth Bockstein $\beta^r(u_1) = u_2$. Let x_i (and y_i) denote the basis for $H_i(K(\mathbb{Z}/2^r;\mathbb{Z}/2)$. The Steenrod operations and the Bockstein on lower homology are given by

$$\begin{array}{rclcrcl} Sq^2_* x_4 & = & x_2 & \qquad & Sq^2_* y_4 & = & y_2 \\ \beta_r(x_4) & = & x_3 & & \beta_r y_4 & = & y_3 \\ \beta_r(x_2) & = & x_1 & & \beta_r y_2 & = & y_1. \end{array}$$

The $\mathbb{Z}/2$-vector space $s^{-1}\tilde{H}_k(X;\mathbb{Z}/2)$ with $k \leq 6$ has a basis given by the table

$$\left(\begin{array}{lllll} k=6 & x_1y_4 & x_2y_3 & x_3y_2 & x_4y_1 \\ 5 & x_1y_3 & x_2y_2 & x_3y_1 & \\ 4 & & & x_1y_2 & x_2y_1 \\ 3 & & & & x_1y_1 \end{array}\right)$$

Let $\alpha \in H_6(X;\mathbb{Z}/2)$ be a spherical class. Then

$$s^{-1}\alpha = \epsilon_1 x_1y_4 + \epsilon_2 x_2y_3 + \epsilon_3 x_3y_2 + \epsilon_4 x_4y_1$$

for some $\epsilon_i \in \mathbb{Z}/2$. Observe that for any spherical class,

$$\beta_s(\alpha) = Sq^t_*(\alpha) = 0$$

for any $s,t \geq 1$. By applying Sq^2_* to α, we have

$$\begin{array}{rcl} 0 & = & Sq^2_*(s^{-1}\alpha) \\ & = & \epsilon_1 Sq^2_*(x_1y_4) + \epsilon_2 Sq^2_*(x_2y_3) + \epsilon_3 Sq^2_*(x_3y_2) + \epsilon_4 Sq^2_*(x_4y_1) \\ & = & \epsilon_1 x_1y_2 + 0 + 0 + \epsilon_4 x_2y_1 \end{array}$$

in $s^{-1}H_4(X;\mathbb{Z}/2)$. Thus

$$\epsilon_1 = \epsilon_4 = 0 \tag{4.2}$$

By applying the Bockstein β_r to α, we have

$$\begin{aligned} 0 &= \beta_r(s^{-1}\alpha) \\ &= \epsilon_1\beta_r(x_1y_4)+\epsilon_2\beta_r(x_2y_3)+\epsilon_3\beta_r(x_3y_2)+\epsilon_4\beta_r(x_4y_1) \\ &= \epsilon_1x_1y_3+\epsilon_2x_1y_3+\epsilon_3x_3y_1+\epsilon_4x_3y_1 \\ &= (\epsilon_1+\epsilon_2)x_1y_3+(\epsilon_3+\epsilon_4)x_3y_1 \end{aligned}$$

and so

$$\epsilon_1+\epsilon_2=\epsilon_3+\epsilon_4=0.$$

Together with equation (4.2), we have $\epsilon_i=0$ for $1\leq i\leq 4$. Thus $\alpha=0$ and hence the result. □

Theorem 4.1. *$\pi_5(\Sigma K(\mathbb{Z}/2,1))\cong\pi_5(\Sigma K(\mathbb{Z}/2,1)\wedge K(\mathbb{Z}/2,1))=\mathbb{Z}/2\oplus\mathbb{Z}/2$.*

Proof. Let $X=\Sigma K(\mathbb{Z}/2,1)\wedge K(\mathbb{Z}/2,1)$. Notice that

$$H_6(X)=\mathbb{Z}/2\oplus\mathbb{Z}/2\cong H_6(X;\mathbb{Z}/2).$$

From diagram (4.1), there is an exact sequence

$$H_6(X)=\mathbb{Z}/2\oplus\mathbb{Z}/2\hookrightarrow\Gamma_5(X)\longrightarrow\pi_4(X)\twoheadrightarrow H_5(X)=\mathbb{Z}/2\oplus\mathbb{Z}/2.$$

By Corollary 3.2,

$$\pi_4(X)\cong\pi_4(\Sigma K(\mathbb{Z}/2,1))\cong\mathbb{Z}/4.$$

From Theorem 2.2(2), there is a short exact sequence

$$\pi_4(X)\otimes\mathbb{Z}/2\oplus\Lambda^2(\pi_3(X))=\mathbb{Z}/2\hookrightarrow\Gamma_5(X)\twoheadrightarrow\operatorname{Tor}(\pi_3(X),\mathbb{Z}/2)=\mathbb{Z}/2.$$

Thus the group $\Gamma_5(X)$ is of order 4. It follows that the monomorphism

$$H_6(X)=\mathbb{Z}/2\oplus\mathbb{Z}/2\hookrightarrow\Gamma_4(X)$$

is an isomorphism and hence the result. □

Lemma 4.3. *Let $r_1,r_2\geq 1$ with $\max\{r_1,r_2\}>1$. Then there is a short exact sequence*

$$\mathbb{Z}/2\oplus\mathbb{Z}/2\hookrightarrow\Gamma_5(\Sigma K(\mathbb{Z}/2^{r_1},1)\wedge K(\mathbb{Z}/2^{r_2},1))\twoheadrightarrow\mathbb{Z}/2.$$

Proof. Let $A=\mathbb{Z}/2^{r_1}\oplus\mathbb{Z}/2^{r_2}$. Let $X=\Sigma K(\mathbb{Z}/2^{r_1},1)\wedge K(\mathbb{Z}/2^{r_2},1)$. Then X is a retract of $\Sigma K(A,1)\wedge K(A,1)$. From Theorem 3.2, $\Gamma_4(X)=\mathbb{Z}/2^{r_1}\otimes\mathbb{Z}/2^{r_2}\otimes\mathbb{Z}/2=\mathbb{Z}/2$ is summand of $\pi_4(X)$ and so

$$\begin{aligned} \pi_4(X) &\cong \Gamma_4(X)\oplus H_4(X) \\ &= \Gamma_4(X)\oplus H_3(K(\mathbb{Z}/2^{r_1},1)\wedge K(\mathbb{Z}/2^{r_2},1)) \\ &\cong \Gamma_4(X)\oplus\operatorname{Tor}(\mathbb{Z}/2^{r_1},\mathbb{Z}/2^{r_2}) \\ &\cong \mathbb{Z}/2\oplus\mathbb{Z}/2^{\min\{r_1,r_2\}}. \end{aligned}$$

The assertion follows from Theorem 2.2(2). □

There is a canonical choice of skeleton $\mathrm{sk}_n(K(\mathbb{Z}/2^r,1))$ with

$$\mathrm{sk}_n(K(\mathbb{Z}/2^r,1)=\mathrm{sk}_{n-1}(K(\mathbb{Z}/2^r,1)\cup e^n.$$

This induces a choice of skeleton

$$\mathrm{sk}_n(\Sigma K(\mathbb{Z}/2^{r_1},1)\wedge K(\mathbb{Z}/2^{r_2},1))=\Sigma\bigcup_{i+j\leq n}\mathrm{sk}_i(K(\mathbb{Z}/2^{r_1},1))\wedge\mathrm{sk}_j(K(\mathbb{Z}/2^{r_2},1)).$$

Lemma 4.4. *Let $r_1,r_2\geq 1$. Let $r=\min\{r_1,r_2\}$. Let $X=\Sigma K(\mathbb{Z}/2^{r_1},1)\wedge K(\mathbb{Z}/2^{r_2},1)$. Then*

(1) $\mathrm{sk}_4(X) \simeq M(\mathbb{Z}/2^r, 3) \vee S^3$.
(2) *If* $r_1 = r_2 = 1$, *then* $\mathrm{sk}_5(X) \simeq S^5 \vee S^5 \vee \Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2$.
(3) *If* $\max\{r_1, r_2\} > 1$, *then* $\mathrm{sk}_5(X) \simeq S^5 \vee S^5 \vee M(\mathbb{Z}/2^r, 3) \vee M(\mathbb{Z}/2^r, 4)$.
(4) $$\Gamma_5(X) \cong \pi_5(\Sigma M(\mathbb{Z}/2^{r_1}, 1) \wedge M(\mathbb{Z}/2^{r_2}, 1)) = \begin{cases} \mathbb{Z}/2 \oplus \mathbb{Z}/2 & \text{if} \quad r_1 = r_2 = 1, \\ \mathbb{Z}/4 \oplus \mathbb{Z}/2 & \text{if} \quad \min\{r_1, r_2\} = 1 \\ & \qquad \text{and } \max\{r_1, r_2\} > 1, \\ \mathbb{Z}/2 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/2 & \text{if} \quad r_1, r_2 > 1. \end{cases}$$

Proof. We may assume that $r_1 \leq r_2$ and so $r = r_1$. Let x_i (y_i) be a basis for $H_i(K(\mathbb{Z}/2^{r_1}, 1); \mathbb{Z}/2)$ ($H_i(K(\mathbb{Z}/2^{r_2}, 1); \mathbb{Z}/2)$), which represents the i-dimensional cell in the space $K(\mathbb{Z}/2^{r_k}, 1)$. Then
$$s^{-1}\tilde{H}_*(\mathrm{sk}_{n+1}(X); \mathbb{Z}/2)$$
has a basis given by $x_i y_j$ with $i + j \leq n$ and $i, j \geq 1$. In particular, $s^{-1}\tilde{H}_*(\mathrm{sk}_4(X); \mathbb{Z}/2)$ has a basis $\{x_1y_1, x_1y_2, x_2y_1\}$ with the Bockstein $\beta_{r_1}(x_2y_1) = x_1y_1$. There is (unique up to homotopy) 2-local 3-cell complex with this homological structure which is given by $S^4 \vee M(\mathbb{Z}/2^r, 3)$. Thus $\mathrm{sk}_4(X) \simeq S^4 \vee M(\mathbb{Z}/2^r, 3)$, which is assertion (1).

(2) and (3). Observe that $s^{-1}\tilde{H}_*(\mathrm{sk}_5(X); \mathbb{Z}/2)$ has a basis $\{x_1y_1, x_1y_2, x_2y_1, x_1y_3, x_2y_2, x_3y_3\}$. Let
$$j\colon \Sigma M(\mathbb{Z}/2^{r_1}, 1) \wedge M(\mathbb{Z}/2^{r_2}, 1) \hookrightarrow \mathrm{sk}_5(X)$$
be the canonical inclusion. For $i = 1, 2$, the composite
$$S^5 \xrightarrow{g} \Sigma^2 K(\mathbb{Z}/2^{r_i}, 1) \cong \Sigma K(\mathbb{Z}/2^{r_i}, 1) \wedge S^1 \hookrightarrow \Sigma K(\mathbb{Z}/2^{r_1}, 1) \wedge K(\mathbb{Z}/2^{r_2}, 1),$$
in which g is map that inducing isomorphism on $H_5(\ ; \mathbb{Z}/2)$ as in Lemma 4.1, induces a map
$$f_i\colon S^5 \longrightarrow \mathrm{sk}_5(X).$$
By inspecting homology, the map
$$(f_1, f_2, j)\colon S^5 \vee S^5 \vee \Sigma M(\mathbb{Z}/2^{r_1}, 1) \wedge M(\mathbb{Z}/2^{r_2}, 1) \longrightarrow \mathrm{sk}_5(X)$$
induces an isomorphism on mod 2 homology and so it is a homotopy equivalent localized at 2. If $\max\{r_1, r_2\} > 1$, then from decomposition 3.7,
$$\Sigma M(\mathbb{Z}/2^{r_1}, 1) \wedge M(\mathbb{Z}/2^{r_2}, 1) \simeq M(\mathbb{Z}/2^r, 3) \vee M(\mathbb{Z}/2^r, 4)$$
and so $\mathrm{sk}_5(X) \simeq S^5 \vee S^5 \vee M(\mathbb{Z}/2^r, 3) \vee M(\mathbb{Z}/2^r, 4)$ in this case. Thus assertions (2) and (3) follow.

(4). **Case I.** $\max\{r_1, r_2\} > 1$. By the definition of the Whitehead's functor Γ,
$$\begin{aligned} \Gamma_5(X) &= \mathrm{Im}(\pi_5(\mathrm{sk}_4(X)) \to \pi_5(\mathrm{sk}_5(X)) \\ &= \mathrm{Im}(\pi_5(S^4 \vee M(\mathbb{Z}/2^r, 3)) \to \pi_5(S^5 \vee S^5 \vee M(\mathbb{Z}/2^r, 3) \vee M(\mathbb{Z}/2^r, 4))) \\ &= \pi_5(M(\mathbb{Z}/2^r, 3) \vee M(\mathbb{Z}/2^r, 4)) \end{aligned}$$
because
$$M(\mathbb{Z}/2^r, 3) \vee M(\mathbb{Z}/2^r, 4))) \simeq \Sigma M(\mathbb{Z}/2^{r_1}, 1) \wedge M(\mathbb{Z}/2^{r_2}, 1) = (S^4 \vee M(\mathbb{Z}/2^r, 3)) \cup e^5.$$
Now it suffices to compute
$$\pi_5(M(\mathbb{Z}/2^r, 3) \vee M(\mathbb{Z}/2^r, 4)) = \pi_5(M(\mathbb{Z}/2^r, 3)) \oplus \pi_5(M(\mathbb{Z}/2^r, 4)).$$
It is straight forward to see that $\pi_5(M(\mathbb{Z}/2^r, 4)) = \mathbb{Z}/2$ represented by the composite
$$S^5 \xrightarrow{\eta} S^4 \hookrightarrow M(\mathbb{Z}/2^r, 4).$$

If $r = \min\{r_1, r_2\} = 1$, then $\pi_5(M(\mathbb{Z}/2^r, 3) = \mathbb{Z}/4$ according to [22, Proposition 5.1].

If $r = \min\{r_1, r_2\} > 1$, we compute $\pi_5(M\mathbb{Z}/2^r, 3)$. Observe that this is in the stable range and so

$$\pi_5(M(\mathbb{Z}/2^r, 3)) \cong \pi_5^s(M(\mathbb{Z}/2^r, 3)).$$

Now we are working in the stable homotopy category. Since $\eta\colon S^5 \to S^4$ is of order 2, there is a map

$$\tilde{\eta}\colon M(\mathbb{Z}/2, 5) \longrightarrow S^4$$

such that $\tilde{\eta}|_{S^5} \simeq \eta$. Since the identity map of $M(\mathbb{Z}/2, 5)$ is of order 4 (see for instance [17, Theorem 4.4]), there is a commutative diagram

$$\begin{tikzcd}
 & S^5 \arrow[r, hook, "j"] & M(\mathbb{Z}/2,5) \arrow[d, "\tilde{\eta}"] \arrow[dll, dashed, "\bar{\eta}"'] & \\
M(\mathbb{Z}/2^r,3) \arrow[rr, "\text{pinch}"] & & S^4 \arrow[r, "{[2^r]}"] & S^4,
\end{tikzcd}$$

where the bottom row is the cofibre sequence. The composite $\bar{\eta}\colon S^5 \to M(\mathbb{Z}/2^r, 3)$ represents an element in $\pi_5^s(M(\mathbb{Z}/2^r, 3))$ that maps down to $\pi_5^s(S^4) = \mathbb{Z}/2(\eta)$. Since the map $j\colon S^5 \to M(\mathbb{Z}/2, 5)$ is of order 2, the composite $\bar{\eta} \circ j$ is of order 2. It follows that

$$\pi_5(M(\mathbb{Z}/2^r, 3)) \cong \pi_5^s(M(\mathbb{Z}/2^r, 3)) \cong \pi_5^s(S^4) \oplus \pi_5^s(S^3) \cong \mathbb{Z}/2 \oplus \mathbb{Z}/2. \tag{4.3}$$

Case II. $r_1 = r_2 = 1$. In this case, similar to the above arguments,

$$\Gamma_5(X) = \Gamma_5(\Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2) = \pi_5(\Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2). \tag{4.4}$$

We compute this homotopy group. Note that

$$\Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2 = \mathrm{sk}_4(X) \cup e^5 = (S^4 \vee M(\mathbb{Z}/2, 3) \cup e^5.$$

There is a cofibre sequence

$$S^4 \xrightarrow{f} S^4 \vee M(\mathbb{Z}/2, 3) \xrightarrow{g} \Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2,$$

where the composite

$$S^4 \xrightarrow{f} S^4 \vee M(\mathbb{Z}/2, 3) \xrightarrow{\text{proj.}} S^4$$

is of degree 2 because

$$Sq^1_*\colon H_5(\Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2; \mathbb{Z}/2) = \mathbb{Z}/2 \longrightarrow H_4(\Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2; \mathbb{Z}/2) = \mathbb{Z}/2 \oplus \mathbb{Z}/2$$

is not zero, and the composite

$$S^4 \xrightarrow{f} S^4 \vee M(\mathbb{Z}/2, 3) \xrightarrow{\text{proj.}} M(\mathbb{Z}/2, 3)$$

is homotopic to the composite

$$S^4 \xrightarrow{\eta} S^3 \xhookrightarrow{j} M(\mathbb{Z}/2, 3)$$

because $\pi_4(M(\mathbb{Z}/2, 3)) = \mathbb{Z}/2$ and

$$Sq^2_*\colon H_5(\Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2; \mathbb{Z}/2) = \mathbb{Z}/2 \longrightarrow H_3(\Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2; \mathbb{Z}/2) = \mathbb{Z}/2$$

is an isomorphism. Since

$$\Gamma_5(\Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2) = \pi_5(\Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2),$$

$$g_*\colon \pi_5(S^4 \vee M(\mathbb{Z}/2, 3)) \longrightarrow \pi_5(\Sigma\mathbb{R}\mathrm{P}^2 \wedge \mathbb{R}\mathrm{P}^2)$$

is an epimorphism. By applying the Hilton-Milnor Theorem,

$$\begin{aligned}
\pi_5(S^4\vee M(\mathbb{Z}/2,3)) &\cong \pi_4(\Omega(S^4\vee M(\mathbb{Z}/2,3)))\\
&\cong \pi_4(\Omega S^4\times \Omega(M(\mathbb{Z}/2,3))\times \Omega\Sigma(\Omega S^4\wedge \Omega M(\mathbb{Z}/2,3)))\\
&\cong \pi_4(\Omega S^4)\oplus \pi_4(\Omega(M(\mathbb{Z}/2,3)))\\
&\cong \pi_5(S^4)\oplus \pi_5(M(\mathbb{Z}/2,3))\\
&\cong \mathbb{Z}/2\oplus \pi_5(M(\mathbb{Z}/2,3)).
\end{aligned}$$

From [22, Proposition 5.1], $\pi_5(M(\mathbb{Z}/2,3))=\mathbb{Z}/4$ generated by the homotopy class of any map $\phi\colon S^5\to M(\mathbb{Z}/2,3)$ such that the composite

$$S^5\to M(\mathbb{Z}/2,3)\to S^4$$

is homotopic to η, the generator for $\pi_5(S^4)=\mathbb{Z}/2$, and, for any such a choice of map ϕ, the element $2[\phi]$ is given by the homotopy class of the composite

$$S^5\xrightarrow{\eta} S^4\xrightarrow{\eta} S^3\xhookrightarrow{j} M(\mathbb{Z}/2,3).$$

From the fact that $g\circ f\simeq *$, the composite

$$\pi_5(S^4)\xrightarrow{f_*}\pi_5(\pi_5(S^4\vee M(\mathbb{Z}/2,3)))\xrightarrow{g_*}\pi_5(\Sigma\mathbb{R}\mathrm{P}^2\wedge\mathbb{R}\mathrm{P}^2)$$

is zero. Observe that

$$f_*(\eta)=2\eta+[j\circ\eta\circ\eta]=2[\phi].$$

Thus $g_*(2[\phi])=0$ and so $\pi_5(\Sigma\mathbb{R}\mathrm{P}^2\wedge\mathbb{R}\mathrm{P}^2)$ is a quotient group $\mathbb{Z}/2\oplus\mathbb{Z}/2$. On the other hand, from Theorem 2.2(2), there is short exact sequence

$$\mathbb{Z}/2\hookrightarrow\Gamma_5(\Sigma\mathbb{R}\mathrm{P}^2\wedge\mathbb{R}\mathrm{P}^2)=\pi_5(\Sigma\mathbb{R}\mathrm{P}^2\wedge\mathbb{R}\mathrm{P}^2)\twoheadrightarrow\mathbb{Z}/2.$$

It follows that $\pi_5(\Sigma\mathbb{R}\mathrm{P}^2\wedge\mathbb{R}\mathrm{P}^2)=\mathbb{Z}/2\oplus\mathbb{Z}/2$. The proof is finished. □

Let $\mathrm{Len}^3(2^r)=\mathrm{sk}_3(K(\mathbb{Z}/2^r,1))$ be the 3-dimensional lens space.

Lemma 4.5. *Let* $r_1,r_2\geq 1$. *Let*

$$\begin{gathered}
X_1=\Sigma M(\mathbb{Z}/2^{r_1},1)\wedge M(\mathbb{Z}/2^{r_2},1),\\
X_2=\Sigma\mathrm{Len}^3(2^{r_1})\wedge\mathrm{Len}^3(2^{r_2}),\\
X=\Sigma K(\mathbb{Z}/2^{r_1},1)\wedge K(\mathbb{Z}/2^{r_2},1).
\end{gathered}$$

Then there is a commutative diagram

$$\begin{tikzcd}
 & \Gamma_5(X_1) \arrow[r, "\cong"] \arrow[d, "\cong"] & \pi_5(X_1) \arrow[d, hook] & \\
 & \Gamma_5(X_2) \arrow[r, hook, "\curvearrowleft"] \arrow[d, "\cong"] & \pi_5(X_2) \arrow[r, two heads, "\curvearrowleft"] \arrow[d, two heads] & \mathbb{Z}/2^{r_1}\oplus\mathbb{Z}/2^{r_2} \arrow[d, two heads] \\
H_6(X) \arrow[r] & \Gamma_5(X) \arrow[r] & \pi_5(X) \arrow[r, two heads] & (\mathbb{Z}/2^{\min\{r_1,r_2\}})^{\oplus 2},
\end{tikzcd}$$

where the rows are exact and the middle a splitting short exact sequence.

Proof. As in the proof in Lemma 4.4, $s^{-1}\tilde{H}_k(X)$ for $k \leq 6$ has a basis

$$\{x_i y_j \mid i + j \leq 6,\ i, j \geq 1\}.$$

Thus

$$\mathrm{sk}_4(X) \subseteq X_1 \subseteq \mathrm{sk}_5(X) \subseteq X_2 \subseteq \mathrm{sk}_7(X)$$

and so the commutative diagram follows, where

$$\Gamma_5(X_1) \cong \Gamma_5(X_2) \cong \Gamma_5(X)$$

are given by Lemma 4.4. Since $\mathrm{sk}_5(X) \subseteq X_2$, $\pi_5(X_2) \to \pi_5(X)$ is onto.

Now we show that the middle row in the diagram splits off. By taking the suspension, there is a commutative diagram of short exact sequences

$$\begin{tikzcd}
\Gamma_5(X_2) \arrow[r, hook] \arrow[d, "\cong"] & \pi_5(X_2) \arrow[r, two heads] \arrow[d] & H_5(X_2) \arrow[d, "\cong"] \\
\Gamma_6(\Sigma X_2) \arrow[r, hook] & \pi_6(\Sigma X_2) \arrow[r, two heads] & H_6(X_2),
\end{tikzcd}$$

where the left column is an isomorphism because

$$\Gamma_5(X_2) \cong \pi_5(X_1) \cong \pi_6(\Sigma X_1) \cong \Gamma_6(\Sigma X_2).$$

Thus

$$\pi_5(X_2) \cong \pi_6(\Sigma X_2) \tag{4.5}$$

by the 5-Lemma.

From Lemma 4.1, there is a map

$$g\colon S^5 \longrightarrow \Sigma^2 K(\mathbb{Z}/2^r, 1)$$

inducing an isomorphism on $H_5(\ ;\mathbb{Z}/2)$. It follows that

$$\Sigma^2\mathrm{Len}^3(2^r) = \Sigma^2\mathrm{sk}_3(K(\mathbb{Z}/2^r, 1)) \simeq S^5 \vee \Sigma^2 M(\mathbb{Z}/2^r, 1) \tag{4.6}$$

and so

$$\begin{aligned}
\Sigma X_2 &= \Sigma^2\mathrm{Len}^3(2^{r_1}) \wedge \mathrm{Len}^3(2^{r_2}) \\
&\simeq (S^5 \vee \Sigma^2 M(\mathbb{Z}/2^{r_1}, 1)) \wedge \mathrm{Len}^3(2^{r_2}) \\
&\simeq \Sigma^5\mathrm{Len}^3(2^{r_2}) \vee \Sigma^2\mathrm{Len}^3(2^{r_2}) \wedge M(\mathbb{Z}/2^{r_1}, 1) \\
&\simeq S^8 \vee M(\mathbb{Z}/2^{r_2}, 6) \vee M(\mathbb{Z}/2^{r_1}, 6) \vee M(\mathbb{Z}/2^{r_2}, 3) \wedge M(\mathbb{Z}/2^{r_1}, 1).
\end{aligned} \tag{4.7}$$

Thus

$$\pi_6(\Sigma X_2) \cong \mathbb{Z}/2^{r_1} \oplus \mathbb{Z}/2^{r_2} \oplus \Gamma_6(\Sigma X_2)$$

and hence the result. □

Theorem 4.2. *Let $r > 1$. Then*

$$\pi_5(\Sigma K(\mathbb{Z}/2^r, 1)) \cong \pi_5(\Sigma K(\mathbb{Z}/2^r, 1) \wedge K(\mathbb{Z}/2^r, 1)) \cong \mathbb{Z}/2 \oplus \mathbb{Z}/2^r \oplus \mathbb{Z}/2^r.$$

Proof. Let $X = \Sigma K(\mathbb{Z}/2^r, 1) \wedge K(\mathbb{Z}/2^r, 1)$. By Lemma 4.5,

$$\pi_5(X) \cong \mathbb{Z}/2^r \oplus \mathbb{Z}/2^r \oplus \mathrm{Im}(\Gamma_5(X) \to \pi_5(X)).$$

From Lemma 4.4,

$$\Gamma_5(X) = \mathbb{Z}/2^{\oplus 3}.$$

By Lemma 4.1, the composite

$$\pi_6(X) \to H_6(X) = \mathbb{Z}/2^r \oplus \mathbb{Z}/2^r \to H_6(X; \mathbb{Z}/2)$$

is zero. Thus

$$H_6(X) = \mathbb{Z}/2^r \oplus \mathbb{Z}/2^r \longrightarrow \Gamma_5(X) = \mathbb{Z}/2 \oplus \mathbb{Z}/2 \oplus \mathbb{Z}/2$$

detects two copies of $\mathbb{Z}/2$-summands in $\Gamma_5(X)$. The proof is finished. □

4.4. **The Group** $\pi_5(\Sigma K(\mathbb{Z}/2^{r_1}, 1) \wedge K(\mathbb{Z}/2^{r_2}, 1))$ **with** $r_1 < r_2$**.** Our computation is given by analyzing the cell structure. Let x_i be a basis for $\tilde{H}_i(K(\mathbb{Z}/2^{r_1}, \mathbb{Z}/2))$ and let y_i be a basis for $\tilde{H}_i(K(\mathbb{Z}/2^{r_2}; \mathbb{Z}/2))$. Then

$$s^{-1}\tilde{H}_k(\Sigma K(\mathbb{Z}/2^{r_1}, 1) \wedge K(\mathbb{Z}/2^{r_2}, 1), \quad k \leq 6,$$

has a basis $\{x_iy_j \mid i + j \leq 6\}$. From the assumption that $r_1 < r_2$, the Steenrod operation and Bockstein are indicated by the following diagram

$$\begin{tikzcd}
k=6 & \boxed{x_2y_3} \arrow[d, "\beta_{r_1}"'] & \boxed{x_1y_4} \arrow[dl, dashed, "\beta_{r_2}"', "0"] \arrow[dd, "Sq^2_*"] & & \boxed{x_3y_2} \arrow[d, dashed, "\beta_{r_2}"'] & \boxed{x_4y_1} \arrow[dl, "\beta_{r_1}"'] \arrow[dd, "Sq^2_*"] \\
k=5 & \boxed{x_1y_3} & & \boxed{x_2y_2} \arrow[dl, "\beta_{r_1}"'] & \boxed{x_3y_1} & \\
k=4 & & \boxed{x_1y_2} & & & \boxed{x_3y_1} \arrow[dl, "\beta_{r_1}"'] \\
k=3 & & & & \boxed{x_1y_1}, &
\end{tikzcd} \tag{4.8}$$

where the dash arrows mean that the next Bockstein β_{r_2}, which comes from $H_*(K(\mathbb{Z}/2^{r_2}, 1))$, does not actually happen in the Bockstein spectral sequence up to this range.

Lemma 4.6. *Let* $r_2 > r_1 \geq 1$ *and let* $X = \Sigma K(\mathbb{Z}/2^{r_1}, 1) \wedge K(\mathbb{Z}/2^{r_2}, 1)$*. Then the suspension*

$$E\colon \pi_5(X) \longrightarrow \pi_6(\Sigma X)$$

is an isomorphism.

Proof. From formula (4.5) together with the fact that $\pi_{n-1}(\mathrm{sk}_n(Y)) \cong \pi_{n-1}(Y)$,

$$\pi_5(\mathrm{sk}_6(\Sigma \mathrm{Len}^3(2^{r_1}) \wedge \mathrm{Len}^3(2^{r_2}))) \longrightarrow \pi_6(\mathrm{sk}_7(\Sigma \mathrm{Len}^3(2^{r_1}) \wedge \mathrm{Len}^3(2^{r_2}))).$$

Notice that

$$\mathrm{sk}_6(X) = \mathrm{sk}_6(\Sigma \mathrm{Len}^3(2^{r_1}) \wedge \mathrm{Len}^3(2^{r_2})) \cup e^6 \cup e^6$$

indicated by the elements x_1y_4 and x_4y_1 in diagram (4.8). Then there is a commutative diagram of right exact sequences

$$\begin{tikzcd}
\pi_5(S^5 \vee S^5) \arrow[r, "f_*"] \arrow[d, "\cong"] & \pi_5(Z) \arrow[r, two heads] \arrow[d, "\cong"] & \pi_5(\mathrm{sk}_6(X)) \arrow[d, "E"] \\
\pi_6(S^6 \vee S^6) \arrow[r, "\Sigma f_*"] & \pi_6(\Sigma Z) \arrow[r, two heads] & \pi_6(\mathrm{sk}_7(\Sigma X)),
\end{tikzcd}$$

where $Z = \mathrm{sk}_6(\Sigma \mathrm{Len}^3(2^{r_1}) \wedge \mathrm{Len}^3(2^{r_2}))$ and $f\colon S^5 \vee S^5 \to Y$ is the attaching map for $\mathrm{sk}_6(X)$. The assertion follows by the 5-lemma. $\square$

Theorem 4.3. *Let $r_2 > r_1 \geq 1$. Then*

$$\pi_5(\Sigma K(\mathbb{Z}/2^{r_1},1) \wedge K(\mathbb{Z}/2^{r_2},1)) = \begin{cases} \mathbb{Z}/2 \oplus \mathbb{Z}/4 & \text{if} \quad r_1 = 1,\ r_2 = 2, \\ \mathbb{Z}/2 \oplus \mathbb{Z}/8 & \text{if} \quad r_1 = 1,\ r_2 \geq 3, \\ \mathbb{Z}/2 \oplus \mathbb{Z}/2^{r_1} \oplus \mathbb{Z}/2^{r_1+1} & \text{if} \quad r_2 > r_1 > 1. \end{cases}$$

Proof. From Lemma 4.6, it suffices to compute $\pi_6(\Sigma^2 K(\mathbb{Z}/2^{r_1},1) \wedge K(\mathbb{Z}/2^{r_2},1))$. Let $X = \mathrm{sk}_7(\Sigma^2 K(\mathbb{Z}/2^{r_1},1) \wedge K(\mathbb{Z}/2^{r_2},1))$. From splitting formula (4.7),

$$\mathrm{sk}_7(\Sigma^2 \mathrm{Len}^3(2^{r_1}) \wedge \mathrm{Len}^3(2^{r_2})) \simeq M(\mathbb{Z}/2^{r_2},6) \vee M(\mathbb{Z}/2^{r_1},6) \vee M(\mathbb{Z}/2^{r_1},4) \vee M(\mathbb{Z}/2^{r_1},5).$$

Let $Y = \mathrm{sk}_7(\Sigma^2 \mathrm{Len}^3(2^{r_1}) \wedge \mathrm{Len}^3(2^{r_2}))$. Then $s^{-2}\tilde{H}_*(Y;\mathbb{Z}/2)$ has a basis listed in diagram (4.8) excluding the elements x_1y_4 and x_4y_1. Let $P^n(2^r) = M(\mathbb{Z}/2^r, n-1)$. The mod homology $\tilde{H}_*(P^n(2^r);\mathbb{Z}/2)$ has a basis u^r_{n-1} and v^r_n with degrees $|u^r_{n-1}| = n-1$, $v^r_n| = n$ and the Bockstein $\beta_r(v^r_n) = u^r_{n-1}$. Since $X = Y \cup e^7 \cup e^7$, there is a cofibre sequence

$$S^6_1 \vee S^6_2 \xrightarrow{f} P^7(2^{r_1}) \vee P^7(2^{r_2}) \vee P^5(2^{r_1}) \vee P^6(2^{r_1}) \xrightarrow{g} X \xrightarrow{q} S^7_1 \vee S^7_2,$$

where

$$g_*(u^{r_1}_6, v^{r_2}_7; u^{r_2}_6, v^{r_2}_7; u^{r_1}_4, v^{r_1}_5; u^{r_1}_5, v^{r_1}_6) = s^2(x_1y_3, x_2y_3; x_3y_1, x_3y_2; x_1y_1, x_2y_1; x_1y_2, x_2y_2)$$

for catching the corresponding elements in $\tilde{H}_*(X;\mathbb{Z}/2)$. Here the map f is the attaching map with $f|_{S^6_1}, f|_{S^6_2}$ corresponding to the homological classes $s^2(x_4y_1)$ and $s^2(x_1y_4)$, respectively. Namely, the induced boundary map $q\colon X \to S^7_1 \vee S^7_2$ has the homological property that $q_*\colon H_7(X;\mathbb{Z}/2) \to H_7(S^7_1 \vee S^7_2)$ is given by

$$q_*(x_2y_3) = q_*(x_3y_2) = 0, \quad q_*(s^2(x_4y_1)) = \iota_1, \quad q_*(s^2(x_1y_4)) = \iota_2,$$

where ι_j is the basis for $H_7(S^7_j;\mathbb{Z}/2)$. For $j = 1,2$, let X_j be the homotopy cofibre of f_j. Then there is a commutative diagram

$$\begin{tikzcd}
S^6_1 \vee S^6_2 \arrow[r, "f"] & P^7(2^{r_1}) \vee P^7(2^{r_2}) \vee P^5(2^{r_1}) \vee P^6(2^{r_1}) \arrow[r, "g"] \arrow[d, equal] & X \arrow[r, "q"] & S^7_1 \vee S^7_2 \\
S^6_j \arrow[u, hook] \arrow[r, "f|_{S^6_j}"] & P^7(2^{r_1}) \vee P^7(2^{r_2}) \vee P^5(2^{r_1}) \vee P^6(2^{r_1}) \arrow[r, "g_j"] & X_j \arrow[u, "\theta_j"] \arrow[r, "q_j"] & S^7_j. \arrow[u, hook]
\end{tikzcd} \tag{4.9}$$

Statement 1. *$\theta_{j*}\colon \tilde{H}_*(X_j;\mathbb{Z}/2) \to \tilde{H}_*(X;\mathbb{Z}/2)$ is a monomorphism. Moreover,*

$$\mathrm{Im}(\theta_{1*}\colon H_7(X_1;\mathbb{Z}/2) \to H_7(X;\mathbb{Z}/2))$$

has the basis given by $\{s^2(x_2y_3), s^2(x_3y_2), s^2(x_4y_1)\}$ *and*

$$\mathrm{Im}(\theta_{2*}\colon H_7(X_2;\mathbb{Z}/2) \to H_7(X;\mathbb{Z}/2))$$

has the basis given by $\{s^2(x_2y_3), s^2(x_3y_2), s^2(x_1y_4)\}$. *Thus a basis for* $\tilde{H}_*(X_j;\mathbb{Z}/2)$ *can be listed in diagram (4.8) by removing one element.* The statement follows immediately by applying mod 2 homology to diagram (4.9), where the only simple computation is given by checking the image of θ_{j*}.

Statement 2. *The composite*

$$\phi_j\colon S^6_j \xrightarrow{f|_{S^6_j}} P^7(2^{r_1}) \vee P^7(2^{r_2}) \vee P^5(2^{r_1}) \vee P^6(2^{r_1}) \xrightarrow{\mathrm{proj.}} P^7(2^{r_1})$$

is null homotopic for $j = 1, 2$.

Consider the commutative diagram of cofibre sequences

$$\begin{tikzcd}
S^6_j \arrow[r, "f|_{S^6_j}"] \arrow[d, equal] & P^7(2^{r_1}) \vee P^7(2^{r_2}) \vee P^5(2^{r_1}) \vee P^6(2^{r_1}) \arrow[r, "g_j"] \arrow[d, "\mathrm{proj.}"'] & X_j \arrow[d, "\delta"] \\
S^6_j \arrow[r] & P^7(2^{r_1}) \arrow[r] & Z.
\end{tikzcd}$$

Then $\dim\tilde{H}_*(Z;\mathbb{Z}/2) = 3$ and $\delta_*\colon H_*(X_j;\mathbb{Z}/2) \to H_*(Z;\mathbb{Z}/2)$ is onto. From diagram (4.8), the Bockstein

$$\beta_t\colon H_7(Z;\mathbb{Z}/2) \to H_6(Z;\mathbb{Z}/2)$$

is 0 for $t < r_1$ with the first non-trivial Bockstein given by β_{r_1} coming from $\beta_{r_1}(x_2y_3) = x_1y_3$ in diagram (4.8). Note that $\pi_6(P^7(2^{r_1})) = \mathbb{Z}/2^{r_1}$ generated by the inclusion $\bar{\iota}\colon S^6 \hookrightarrow P^7(2^{r_1})$. Then the homotopy class

$$[\phi_j] = k\bar{\iota}$$

for some $k \in \mathbb{Z}$. If $k \equiv 1 \mod 2$, then $\dim\tilde{H}_*(Z;\mathbb{Z}/2) = 1$ which contradicts to that $\dim\tilde{H}_*(Z;\mathbb{Z}/2) = 3$. Thus k must be divisible by 2. Let $k = 2^tk'$ with $k' \equiv 1 \mod 2$ for some $t \geq 1$. If $t < r_1$, then there is a nontrivial Bockstein β_t on $\tilde{H}_*(Z;\mathbb{Z}/2)$ which is impossible from the above. Hence $t \geq r_1$ and so $[\phi_j] = 0$ in $\pi_6(P^7(2^{r_1})) = \mathbb{Z}/2^{r_1}$. Statement 2 follows.

Statement 3. *The composite*

$$\psi\colon S^6_1 \xrightarrow{f|_{S^6_1}} P^7(2^{r_1}) \vee P^7(2^{r_2}) \vee P^5(2^{r_1}) \vee P^6(2^{r_1}) \xrightarrow{\mathrm{proj.}} P^6(2^{r_1})$$

is null homotopic.

Consider the commutative diagram of cofibre sequences

$$\begin{tikzcd}
S^6_1 \arrow[r, "f|_{S^6_1}"] \arrow[d, equal] & P^7(2^{r_1}) \vee P^7(2^{r_2}) \vee P^5(2^{r_1}) \vee P^6(2^{r_1}) \arrow[r, "g_1"] \arrow[d, "p\,\mathrm{proj.}"'] & X_1 \arrow[d, "\delta"] \\
S^6_1 \arrow[r] & P^6(2^{r_1}) \arrow[r, "g'"] & W.
\end{tikzcd}$$

Then $\dim\tilde{H}_*(W;\mathbb{Z}/2)=3$ and $\delta_*\colon \tilde{H}_(X_1;\mathbb{Z}/2)\to\tilde{H}_*(W;\mathbb{Z}/2)$ is onto. Moreover, $H_7(W;\mathbb{Z}/2)$ has a basis given by $\delta_*(s^2(x_4y_1))$. By Statement 1, a basis for $\tilde{H}_*(X_1;\mathbb{Z}/2)$ is listed in diagram (4.8) by removing x_1y_4. The canonical projection

$$p\colon P^7(2^{r_1})\vee P^7(2^{r_2})\vee P^5(2^{r_1})\vee P^6(2^{r_1})\longrightarrow P^6(2^{r_1})$$

has the property that $p_*(u_5^{r_1})=u_5^{r_1}$, $p_*(v_6^{r_1})=v_6^{r_1}$ and $p_*(x)=0$ for x to the other elements in the basis for $\tilde{H}_*(P^7(2^{r_1})\vee P^7(2^{r_2})\vee P^5(2^{r_1})\vee P^6(2^{r_1});\mathbb{Z}/2)$. In particular, $p_*(v_5^{r_1})=0$. Note that

$$\begin{aligned}Sq^2_*\delta_*(s^2(x_4y_1)) &= \delta_*(Sq^2_*(s^2(x_4y_1)))\\ &= \delta_*(s^2(x_2y_1))\\ &= \delta_*(g_{1*}(v_5^{r_1})\\ &= g'_*\circ p_*(v_5^{r_1})\\ &= 0.\end{aligned}$$

If follows that

$$Sq^2_*\colon H_7(W;\mathbb{Z}/2)\to H_5(W;\mathbb{Z}/2)$$

is zero. From the exact sequence

$$\pi_6(S^5)=\mathbb{Z}/2\xrightarrow[0]{2^{r_1}}\pi_6(S^5)=\mathbb{Z}/2\longrightarrow\pi_6(P^6(2^{r_1})\longrightarrow\pi_5(S^5)=\mathbb{Z}\xrightarrow{2^{r_1}}\mathbb{Z},$$

we have

$$\pi_6(P^6(2^{r_1}))=\mathbb{Z}/2 \tag{4.10}$$

generated by the composite

$$\bar{\eta}\colon S^6\xrightarrow{\eta}S^5\hookrightarrow P^6(2^{r_1}).$$

Thus the homotopy class $[\psi]=0$ or $\bar{\eta}$. If $[\psi]=\bar{\eta}$, then $Sq^2\colon H_7(W;\mathbb{Z}/2)\to H_5(W;\mathbb{Z}/2)$ is not zero, which is impossible from the above. Hence $[\psi]=0$. This finishes the proof for Statement 3.

Statement 4. *There is a homotopy decomposition*

$$X\simeq P^7(2^{r_1})\vee T_1\vee T_2,$$

where $\tilde{H}_*(T_1;\mathbb{Z}/2)$ and $\tilde{H}_*(T_2;\mathbb{Z}/2)$ have basis listed by the middle and the right modules in diagram (4.6), respectively.

From Statements 2 and 3, the attaching map $f|_{S^6_1}$ maps into the subspace $P^7(2^{r_2})\vee P^5(2^{r_1})$ up to homotopy because, in the range of π_6, we have

$$\pi_6(P^7(2^{r_1})\vee P^7(2^{r_2})\vee P^5(2^{r_1})\vee P^6(2^{r_1}))\cong\pi_6(P^7(2^{r_1}))\oplus\pi_6(P^7(2^{r_2}))\oplus\pi_6(P^5(2^{r_1}))\oplus\pi_6(P^6(2^{r_1})).$$

Thus there is a homotopy commutative diagram of cofibre sequences

$$\begin{tikzcd}
S^6_1 \arrow[r, "f'"] \arrow[d, hook] & P^7(2^{r_2}) \vee P^5(2^{r_1}) \arrow[r] \arrow[d, hook] & T_2 \arrow[d, dashed, "i_1"] \\
S^6_1 \vee S^6_2 \arrow[r, "f"] \arrow[d, "\text{proj.}"] & P^7(2^{r_1}) \vee P^7(2^{r_2}) \vee P^5(2^{r_1}) \vee P^6(2^{r_1}) \arrow[r, "g"] \arrow[d, "\text{proj.}"] & X \arrow[d, dashed, "q_1"] \\
S^6_1 \arrow[r, "f'"] & P^7(2^{r_2}) \vee P^5(2^{r_1}) \arrow[r] & T_2.
\end{tikzcd} \tag{4.11}$$

From Statement 2, there is a homotopy commutative diagram of cofibre sequences

$$\begin{tikzcd}
S^6_1 \vee S^6_2 \arrow[r, "f"] \arrow[d] & P^7(2^{r_1}) \vee P^7(2^{r_2}) \vee P^5(2^{r_1}) \vee P^6(2^{r_1}) \arrow[r, "g"] \arrow[d, "\text{proj.}"] & X \arrow[d, dashed, "q_2"] \\
\ast \arrow[r] & P^7(2^{r_1}) \arrow[r, equal] & P^7(2^{r_1}).
\end{tikzcd}$$

Now the composite

$$P^7(2^{r_1}) \vee T_2 \xrightarrow{(g|_{P^7(2^{r_1})}, i_1)} X \xrightarrow{(q_2, q_1)} P^7(2^{2^{r_1}}) \vee T_2$$

is a homotopy equivalence by inspecting the homology and hence the statement.

Computation of the Homotopy Group: From Statement 4, we have

$$\pi_6(X) \cong \pi_6(P^7(2^{r_1}) \vee T_1 \vee T_2) \cong \pi_6(P^7(2^{r_1})) \oplus \pi_6(T_1) \oplus \pi_6(T_2) \cong \mathbb{Z}/2^{r_1} \oplus \pi_6(T_1) \oplus \pi_6(T_2).$$

For computing $\pi_6(T_1)$, since $T_1 = P^6(2^{r_1}) \cup e^7$, there is a right exact sequence

$$\pi_6(S^6) = \mathbb{Z} \longrightarrow \pi_6(P^6(2^{r_1})) = \mathbb{Z}/2 \twoheadrightarrow \pi_6(T_1),$$

where $\pi_6(P^6(2^{r_1})) = \mathbb{Z}/2$ is given in formula (4.10). From diagram (4.6),

$$Sq^2_*\colon H_7(T_1; \mathbb{Z}/2) \longrightarrow H_5(T_1; \mathbb{Z}/2)$$

is an isomorphism and so the attaching map $S^6 \to P^6(2^{r_1})$ of T_1 is non-trivial. It follows that $\pi_6(T_1) = 0$.

Now we compute $\pi_6(T_2)$. From diagram 4.11, there is a right exact sequence

$$\pi_6(S^6) = \mathbb{Z}/ \xrightarrow{f'_*} \pi_6(P^7(2^{r_2}) \vee P^5(2^{r_1})) = \pi_6(P^7(2^{r_2})) \oplus \pi_6(P^5(2^{r_1})) \twoheadrightarrow \pi_6(T_2).$$

Note that a basis for $\tilde{H}_*(T_2)$ can be listed in the right module of diagram (4.6). The composite

$$\pi_6(S^6) = \mathbb{Z} \xrightarrow{f'_*} \pi_6(P^7(2^{r_2})) \oplus \pi_6(P^5(2^{r_1})) \xrightarrow{\text{proj.}} \pi_6(P^7(2^{r_2})) = \mathbb{Z}/2^{r_2}$$

is of degree 2^{r_1} because of the existence of the Bockstein β_{r_1}. Moveover the composite

$$S^6 \xrightarrow{f'} P^7(2^{r_2}) \vee P^5(2^{r_1}) \xrightarrow{\text{proj.}} P^5(2^{r_1}) \xrightarrow{\text{pinch}} S^5 \tag{4.12}$$

is homotopic to η because of the existence of the Steenrod operation Sq^2_*.

Case I. $r_1 = 1$. According to [22, Proposition 5.1], $\pi_6(P^5(2)) = \mathbb{Z}/4$ generated by the homotopy class of any may $S^6 \to P^5(2)$ such that the composite $S^6 \to P^5(2) \to S^5$ is η. It follows that there is a right exact sequence

$$\mathbb{Z} \xrightarrow{f'_*=(2^{r_1},\lambda)} \mathbb{Z}/2^{r_2} \oplus \mathbb{Z}/4 \twoheadrightarrow \pi_6(T_2),$$

where $\lambda\colon \mathbb{Z} \to \mathbb{Z}/4$ is an epimorphism. Thus

$$\pi_6(T_2) = \begin{cases} \mathbb{Z}/4 & \text{if} \quad r_2 = 2, \\ \mathbb{Z}/8 & \text{if} \quad r_2 \geq 3. \end{cases} \tag{4.13}$$

Case II. $r_1 > 1$. From formula 4.3, we have $\pi_6(P^5(2^{r_1})) = \mathbb{Z}/2 \oplus \mathbb{Z}/2$. Since the composite in (4.12) is essential, the composite

$$\pi_6(S^6) = \mathbb{Z} \xrightarrow{f'_*} \pi_6(P^7(2^{r_2})) \oplus \pi_6(P^5(2^{r_1})) \xrightarrow{\text{proj.}} \pi_6(P^5(2^{r_1})) = \mathbb{Z}/2 \oplus \mathbb{Z}/2$$

is nontrivial and so there is right exact sequence

$$\mathbb{Z} \xrightarrow{(2^{r_1},\lambda)} \mathbb{Z}/2^{r_2} \oplus (\mathbb{Z}/2 \oplus \mathbb{Z}/2) \twoheadrightarrow \pi_6(T_2)$$

with $\lambda\colon \mathbb{Z} \to \mathbb{Z}/2 \oplus \mathbb{Z}/2$ nontrivial. It follows that

$$\pi_6(T_2) = \mathbb{Z}/2^{r_1+1} \oplus \mathbb{Z}/2 \text{ for } r_1 > 1. \tag{4.14}$$

The proof is finished now. □

4.5. **The Group** $\pi_5(\Sigma^2 K(\mathbb{Z}/2^r, 1))$**.** We use the spectral sequence induced from Carlsson's construction for computing this group. Let A be an abelian group and

$$0 \to A_1 \xrightarrow{\delta} A_0 \to A \to 0$$

a two-step flat resolution of A, i.e. A_0 is a free abelian group. The diagram (1.1) implies that there is a natural isomorphism

$$\pi_4(\Sigma^2 K(A, 1)) \simeq A \tilde{\otimes} A,$$

where $\tilde{\otimes}^2$:

$$\tilde{\otimes}^2(A) = A \tilde{\otimes} A := A \otimes A / (a \otimes b + b \otimes a, \ a, b \in A).$$

Given a free abelian group $\bar{A}$, theorem 2.2 (2) implies the following natural exact sequence:

$$\begin{tikzcd}
\Gamma_5(\Sigma^2 K(\bar{A},1)) \arrow[r, hook] \arrow[d, equal] & \pi_5(\Sigma^2 K(\bar{A},1)) \arrow[r, two heads] \arrow[d, equal] & H_5(\Sigma^2 K(\bar{A},1)) \arrow[d, equal] \\
\bar{A} \tilde{\otimes} \bar{A} \otimes \mathbb{Z}/2 \oplus \Lambda^2(\bar{A}) \arrow[r, hook] & \pi_5(\Sigma^2 K(\bar{A},1)) \arrow[r, two heads] & \Lambda^3(\bar{A})
\end{tikzcd}$$

The spectral sequence (2.9) for $n=2$, gives the following diagram of exact sequences:

$$\begin{tikzcd}
L_1\Lambda^3(A) \arrow[d] & & \\
A\widetilde{\otimes}A\otimes\mathbb{Z}/2\oplus\Lambda^2(A) \arrow[r, hook] \arrow[d] & \pi_5(\Sigma^2K(A,1)) \arrow[r, two heads] \arrow[d, equal] & L_1\widetilde{\otimes}^2(A) \arrow[d, equal] \\
\pi_0(\pi_5(\Sigma^2K(N^{-1}(A_1\xrightarrow{\delta}A_0),1))) \arrow[r, hook] \arrow[d, two heads] & \pi_5(\Sigma^2K(A,1)) \arrow[r, two heads] & \pi_1(\pi_4\Sigma^2K(N^{-1}(A_1\xrightarrow{\delta}A_0),1)) \\
\Lambda^3(A) & &
\end{tikzcd}$$

(4.15)

Consider the first derived functor of the functor $\widetilde{\otimes}^2$. The short exact sequence

$$LSP^2(A)\to L\otimes^2(A)\to L\widetilde{\otimes}^2(A)$$

in the derived category has the following model:

$$\begin{tikzcd}
\Lambda^2(A_1) \arrow[r, hook, "\delta_2"] \arrow[d, hook] & A_1\otimes A_0 \arrow[r, "\delta_1"] \arrow[d, hook] & SP^2(A_0) \arrow[d, hook] \\
A_1\otimes A_1 \arrow[r, hook, "\delta_2'"] \arrow[d, two heads] & (A_1\otimes A_0)\oplus(A_0\otimes A_1) \arrow[r, "\delta_1'"] \arrow[d, two heads] & A_0\otimes A_0 \arrow[d, two heads] \\
SP^2(A_1) \arrow[r, hook, "\delta_2''"] & A_1\otimes A_0 \arrow[r, "\delta_1''"] & A_0\widetilde{\otimes}A_0
\end{tikzcd}$$

with

$$\begin{aligned}
&\delta_2(a_1\wedge a_1')=a_1\otimes\delta(a_1')-a_1'\otimes\delta(a_1)\\
&\delta_1(a_1\otimes a_0)=a_0\delta(a_1)\\
&\delta_2'(a_1\otimes a_1')=(a_1\otimes\delta(a_1'),-a_1'\otimes\delta(a_1))\\
&\delta_1'(a_1\otimes a_0,a_1'\otimes a_0')=\delta(a_1)\otimes a_0+\delta(a_1')\otimes a_0'\\
&\delta_2''(a_1a_1')=a_1\otimes\delta(a_1')+a_1'\otimes\delta(a_1)\\
&\delta_1''(a_1\otimes a_0)=\partial(a_1)\widetilde{\otimes}a_0
\end{aligned}$$

for $a_0,a_0'\in A_0,\ a_1,a_1'\in A_1$. For $n\geq 2$, looking at the resolution $\mathbb{Z}\xrightarrow{n}\mathbb{Z}$ of the cyclic group $\mathbb{Z}/n$, we obtain the following representative of the element $L\widetilde{\otimes}^2(\mathbb{Z}/n)$ in the derived category:

$$\mathbb{Z}\xrightarrow{2n}\mathbb{Z}\xrightarrow{n}\mathbb{Z}/2$$

In particular,

$$L_1\widetilde{\otimes}^2(\mathbb{Z}/2^k) = \mathbb{Z}/2^{k+1},\ k \geq 1. \tag{4.16}$$

Here $L_1\widetilde{\otimes}^2$ denotes the first derived functor of $\widetilde{\otimes}^2$ (see 2.2).

We will use the following:

Lemma 4.7. *(Lemma 2.1 from [21]) Let G_* be a simplicial group and let $n \geq 0$. Suppose that $\pi_0(G_*)$ acts trivially on $\pi_n(G_*)$. Then the homotopy group $\pi_n(G_*)$ is contained in the center of $G_n/\mathcal{B}G_n$, where $\mathcal{B}G_n$ is the nth simplicial boundary subgroup of G_n.* □

Theorem 4.4. *The homotopy group*

$$\pi_5(\Sigma^2 K(\mathbb{Z}/2^r, 1)) = \begin{cases} \mathbb{Z}/8 & \textit{if} \quad r = 1 \\ \mathbb{Z}/2^{r+1} \oplus \mathbb{Z}/2 & \textit{if} \quad r > 1. \end{cases}$$

Proof. **Case 1:** $r = 1$**.** The natural epimorphism $\mathbb{Z} \to \mathbb{Z}/2$ induces the homomorphisms

$$\pi_n(S^3) = \pi_n(\Sigma^2 K(\mathbb{Z}, 1)) \to \pi_n(\Sigma^2 K(\mathbb{Z}/2, 1)) = \pi_n(\Sigma^2\mathbb{R}P^\infty),\ n \geq 1.$$

The diagram (4.15) together with (4.16) implies the following short exact sequences:

$$\begin{tikzcd}
\mathbb{Z}/2 \arrow[r, equal] \arrow[d, equal] & \pi_5(S^3) \arrow[d] & \\
\mathbb{Z}/2 \arrow[r, hook] & \pi_5(\Sigma^2\mathbb{R}P^\infty) \arrow[r, two heads] & \mathbb{Z}/4
\end{tikzcd} \tag{4.17}$$

Consider this map simplicially, at the level of the natural map between the Carlsson constructions $F(S^2) = F^{\mathbb{Z}}(S^2) \to F^{\mathbb{Z}/2}(S^2)$:

$$\begin{tikzcd}
F(S^2)_4 \arrow[r, shift left=3] \arrow[r, shift left=1, "\cdots"] & F(S^2)_3 \arrow[l, shift left=1, "\cdots"] \arrow[l, shift left=3] \arrow[r, shift left=2] \arrow[r] \arrow[r, shift right=2] & \mathbb{Z} \arrow[l, shift left=1] \arrow[l, shift left=3] \\
F(S^2)_4 \arrow[u, phantom] & & \\[-2em]
\end{tikzcd}$$

$$\begin{tikzcd}
F(S^2)_4 \arrow[d] \arrow[r, shift left=3] \arrow[r, shift left=1, "\cdots"] & F(S^2)_3 \arrow[d] \arrow[l, shift left=1, "\cdots"] \arrow[l, shift left=3] \arrow[r, shift left=2] \arrow[r] \arrow[r, shift right=2] & \mathbb{Z} \arrow[d] \arrow[l, shift left=1] \arrow[l, shift left=3] \\
F^{\mathbb{Z}/2}(S^2)_4 \arrow[r, shift left=3] \arrow[r, shift left=1, "\cdots"] & F^{\mathbb{Z}/2}(S^2)_3 \arrow[l, shift left=1, "\cdots"] \arrow[l, shift left=3] \arrow[r, shift left=2] \arrow[r] \arrow[r, shift right=2] & F^{\mathbb{Z}/2}(S^2)_2 \arrow[l, shift left=1] \arrow[l, shift left=3]
\end{tikzcd}$$

Here $F^{\mathbb{Z}/2}(S^2)_k$ is the free product of $\binom{k}{2}$ copies of $\mathbb{Z}/2$. In particular

$$F^{\mathbb{Z}/2}(S^2)_4 = \langle s_j s_i(\sigma)\ 0 \leq i < j \leq 3 |\ (s_j s_i(\sigma))^2 = 1\rangle$$

Using the description of the element (2.7), we see that the simplicial cycle which defines the image of $\pi_5(S^3)$ in $\pi_5(\Sigma^2\mathbb{R}P^\infty)$ can be chosen of the form

$$[[s_2s_1(\sigma), s_1s_0(\sigma)], [s_2s_1(\sigma), s_2s_0(\sigma)]] \in F^{\mathbb{Z}/2}(S^2)_4$$

With the help of lemma 4.7, we have

$$\begin{aligned}
[[s_2s_1(\sigma), s_1s_0(\sigma)], [s_2s_1(\sigma), s_2s_0(\sigma)]] = [[(s_2s_1(\sigma), s_1s_0(\sigma)], (s_2s_1(\sigma)s_2s_0(\sigma))^2] \equiv \\
[[s_2s_1(\sigma), s_1s_0(\sigma)], (s_2s_1(\sigma), s_2s_0(\sigma))]^2 \quad \text{mod } \mathcal{B}^{\mathbb{Z}/2}(S^2)_4
\end{aligned}$$

since $[[s_2s_1(\sigma), s_1s_0(\sigma)], s_2s_1(\sigma)s_2s_0(\sigma)]$ is a cycle in $F^{\mathbb{Z}/2}(S^2)$. That is, the image of the element $\pi_5(S^3)$ is divisible by 2 in $\pi_5(\Sigma^2\mathbb{R}P^\infty)$. The diagram (4.17) implies the result.

Case 2: $r > 1$**.** Now the diagram (4.15) together with (4.16) implies the following short exact sequence

$$0 \to \mathbb{Z}/2 \to \pi_5(\Sigma^2 K(\mathbb{Z}/2^r, 1)) \to \mathbb{Z}/2^{r+1} \to 0 \tag{4.18}$$

Therefore, $\pi_5(\Sigma^2 K(\mathbb{Z}/2^r, 1))$ is either $\mathbb{Z}/2^{r+2}$ or $\mathbb{Z}/2^{r+1} \oplus \mathbb{Z}/2$.

By theorem 2.2 (2), the Whitehead exact sequence for $\Sigma^2 K(A, 1)$ has the following form:

$$\begin{tikzcd}
 & A\widetilde{\otimes}A \otimes \mathbb{Z}/2 \oplus \Lambda^2(A) \arrow[d, hook] & & \\
H_4(A) \arrow[r] & \Gamma_5(\Sigma^2 K(A,1)) \arrow[r] \arrow[d, two heads] & \pi_5(\Sigma^2 K(A,1)) \arrow[r] & H_3(A) \\
 & Tor(A, \mathbb{Z}/2) & &
\end{tikzcd} \tag{4.19}$$

For $A = \mathbb{Z}/2^r$ it is of the following form:

$$\begin{tikzcd}
\mathbb{Z}/2 \arrow[d, hook] & & \\
\Gamma_5(\Sigma^2 K(\mathbb{Z}/2^r,1)) \arrow[r, hook] \arrow[d, two heads] & \pi_5(\Sigma^2 K(\mathbb{Z}/2^r,1)) \arrow[r, two heads] & \mathbb{Z}/2^r \\
\mathbb{Z}/2 & &
\end{tikzcd} \tag{4.20}$$

The natural projection $\mathbb{Z}/2^r \twoheadrightarrow \mathbb{Z}/2$ induces the map

$$\begin{tikzcd}
\mathbb{Z}/2 \arrow[r, "\simeq"] \arrow[d, hook] & \mathbb{Z}/2 \arrow[d, hook] \\
\Gamma_5(\Sigma^2 K(\mathbb{Z}/2^r,1)) \arrow[r] \arrow[d, two heads] & \Gamma_5(\Sigma^2 \mathbb{R}P^\infty) \arrow[d, two heads] \\
\mathbb{Z}/2 \arrow[r, "0"] & \mathbb{Z}/2
\end{tikzcd} \tag{4.21}$$

where the lower map is zero since the induced map $Tor(\mathbb{Z}/2^r, \mathbb{Z}/2) \to Tor(\mathbb{Z}/2, \mathbb{Z}/2)$ is zero. The fact that $\pi_5(\Sigma^2 \mathbb{R}P^\infty) = \mathbb{Z}/8$ together with diagram (4.19) implies that $\Gamma_5(\Sigma^2 \mathbb{R}P^\infty) = \mathbb{Z}/4$. Hence $\Gamma_5(\Sigma^2 K(\mathbb{Z}/2^r, 1)) = \mathbb{Z}/2 \oplus \mathbb{Z}/2$, since there is no any endomorphism $\mathbb{Z}/4 \to \mathbb{Z}/4$ with zero map on quotients $\mathbb{Z}/2 \to \mathbb{Z}/2$ (as in diagram (4.21)). The diagram (4.20) and exact sequence (4.18) implies that $\pi_5(\Sigma^2 K(\mathbb{Z}/2^r, 1)) = \mathbb{Z}/2^{r+1} \oplus \mathbb{Z}/2$. □

4.6. **Applications.**

Proposition 4.7. *The group* $\Gamma_5(\Sigma\mathbb{R}P^\infty)=\mathbb{Z}/2\oplus\mathbb{Z}/2\oplus\mathbb{Z}/2$.

Proof. Consider the Whitehead exact sequence

$$\pi_6(\Sigma\mathbb{R}\mathrm{P}^\infty)\xrightarrow[0]{h} H_6(\Sigma\mathbb{R}\mathrm{P}^\infty)=\mathbb{Z}/2\longrightarrow\Gamma_5(\Sigma\mathbb{R}\mathrm{P}^\infty)\longrightarrow$$
$$\pi_5(\Sigma\mathbb{R}\mathrm{P}^\infty)=\mathbb{Z}/2\oplus\mathbb{Z}/2\longrightarrow H_5(\Sigma\mathbb{R}\mathrm{P}^\infty)=0,$$

where $\pi_5(\Sigma\mathbb{R}\mathrm{P}^\infty)=\mathbb{Z}/2\oplus\mathbb{Z}/2$ by Theorem 4.1 and the Hurewicz homomorphism

$$h_6\colon\pi_6(\Sigma\mathbb{R}\mathrm{P}^\infty)\to H_6(\Sigma\mathbb{R}\mathrm{P}^\infty)$$

is zero because, otherwise, it would induces a splitting of $\Sigma\mathbb{R}\mathrm{P}^5$ which is impossible by inspecting the Steenrod operation on mod 2 homology. Thus the order of $\Gamma_5(\Sigma\mathbb{R}\mathrm{P}^\infty)$ is 8. We have to determine the group $\Gamma_5(\Sigma\mathbb{R}\mathrm{P}^\infty)$. By the definition,

$$\Gamma_5(\Sigma\mathbb{R}\mathrm{P}^\infty)=\mathrm{Im}(\pi_5(\Sigma\mathbb{R}\mathrm{P}^3)\to\pi_5(\Sigma\mathbb{R}\mathrm{P}^4)).$$

Thus the inclusion $\Sigma\mathbb{R}\mathrm{P}^3\to\Sigma\mathbb{R}\mathrm{P}^\infty$ induces an epimorphism

$$\Gamma_5(\Sigma\mathbb{R}\mathrm{P}^3)\twoheadrightarrow\Gamma_5(\Sigma\mathbb{R}\mathrm{P}^\infty).$$

Note that $\mathbb{R}\mathrm{P}^3=SO(3)$ and so, by the Hopf fibration,

$$\begin{aligned}\pi_5(\Sigma SO(3)) &\cong \pi_5(BSO(3))\oplus\pi_5(\Sigma SO(3)\wedge SO(3))\\ &\cong \pi_4(SO(3))\oplus\pi_5(\Sigma\mathbb{R}\mathrm{P}^3\wedge\mathbb{R}\mathrm{P}^3)\\ &\cong \mathbb{Z}/2\oplus\pi_5(\Sigma\mathbb{R}\mathrm{P}^3\wedge\mathbb{R}\mathrm{P}^3).\end{aligned}$$

From Lemmas 4.4 and 4.5,

$$\begin{aligned}\pi_5(\Sigma\mathbb{R}\mathrm{P}^3\wedge\mathbb{R}\mathrm{P}^3) &\cong \Gamma_5(\Sigma\mathbb{R}\mathrm{P}^3\wedge\mathbb{R}\mathrm{P}^3)\oplus\mathbb{Z}/2\oplus\mathbb{Z}/2\\ &\cong \pi_5(\Sigma\mathbb{R}\mathrm{P}^2\wedge\mathbb{R}\mathrm{P}^2)\oplus\mathbb{Z}/2\oplus\mathbb{Z}/2\\ &\cong \mathbb{Z}/2\oplus\mathbb{Z}/2\oplus\mathbb{Z}/2\oplus\mathbb{Z}/2.\end{aligned}$$

It follows that its quotient $\Gamma_5(\Sigma\mathbb{R}\mathrm{P}^\infty)$ must be an elementary 2-group and so hence the result. □

Proposition 4.8. *For the suspended projective spaces,*

$$\pi_5(\Sigma\mathbb{R}\mathrm{P}^n)=\begin{cases}\mathbb{Z}/2 & \text{if } n=1,\\ \mathbb{Z}/2^{\oplus 3} & \text{if } n=2,\\ \mathbb{Z}/2^{\oplus 5} & \text{if } n=3,\\ \mathbb{Z}/2^{\oplus 3} & \text{if } n=4,\\ \mathbb{Z}/2^{\oplus 2} & \text{if } 3\le n\le\infty.\end{cases}$$

Proof. When $n=1$, $\pi_5(S^2)=\mathbb{Z}/2$ from Toda's table[16]. When $n=2$, $\pi_5(\Sigma\mathbb{R}\mathrm{P}^2)=\mathbb{Z}/2^{\oplus 3}$ is given in [22, Theorem 6.36]. When $n=3$, $\pi_5(\Sigma\mathbb{R}\mathrm{P}^3)$ has been computed in Proposition 4.7. For $n\ge 4$, since $\mathrm{sk}_6(\Sigma\mathbb{R}\mathrm{P}^\infty)=\Sigma\mathbb{R}\mathrm{P}^5$,

$$\pi_5(\Sigma\mathbb{R}\mathrm{P}^n)\cong\pi_5(\Sigma\mathbb{R}\mathrm{P}^\infty)=\mathbb{Z}/2\oplus\mathbb{Z}/2$$

by Theorem 4.2. The remaining case is $\pi_5(\Sigma\mathbb{R}\mathrm{P}^4)$. Let F be the homotopy fibre of the pinch map $\Sigma\mathbb{R}\mathrm{P}^6\longrightarrow\Sigma\mathbb{R}\mathrm{P}^6/\mathbb{R}\mathrm{P}^4=M(\mathbb{Z}/2,6)$. By inspecting the Serre spectral sequence to the fibre sequence

$$\Omega M(\mathbb{Z}/2,6)\longrightarrow F\longrightarrow\Omega\Sigma\mathbb{R}\mathrm{P}^6,$$

the canonical injection $j\colon \Sigma\mathbb{R}\mathrm{P}^4 \to F$ induces an isomorphism on $H_k(\ ;\mathbb{Z}/2)$ for $k \leq 6$ and so

$$j_*\colon \pi_k(\Sigma\mathbb{R}\mathrm{P}^4) \longrightarrow \pi_k(F)$$

is an isomorphism for $k \leq 5$. In particular, $\pi_5(\Sigma\mathbb{R}\mathrm{P}^4) \cong \pi_5(F)$. From the exact sequence

$$\pi_5(\Omega M(\mathbb{Z}/2,6)) = \mathbb{Z}/2 \longrightarrow \pi_5(F) \longrightarrow \pi_5(\Sigma\mathbb{R}\mathrm{P}^6) = \mathbb{Z}/2 \oplus \mathbb{Z}/2,$$

the group $\pi_5(F)$ is of order at most 8 and so is $\pi_5(\Sigma\mathbb{R}\mathrm{P}^4)$. From Proposition 4.7,

$$\Gamma_5(\Sigma\mathbb{R}\mathrm{P}^\infty) = \mathrm{Im}(\pi_5(\Sigma\mathbb{R}\mathrm{P}^3) \to \pi_5(\Sigma\mathbb{R}\mathrm{P}^4)) = \mathbb{Z}/2^{\oplus 3}.$$

It follows that $\pi_5(\Sigma\mathbb{R}\mathrm{P}^4) = \mathbb{Z}/2^{\oplus 3}$ and hence the result. □

Proposition 4.9. $\pi_5(\Sigma K(\Sigma_3,1)) \simeq \mathbb{Z}/2 \oplus \mathbb{Z}/2$.

Proof. This follows from the analysis of the map between the Whitehead exact sequences (2.2) induced by the natural map $\mathbb{Z}/2 \hookrightarrow \Sigma_3$:

$$\begin{tikzcd}
H_5(\mathbb{Z}/2) \arrow[r] \arrow[d, equal] & \Gamma_5(\Sigma\mathbb{R}P^\infty) \arrow[r] \arrow[d, equal] & \pi_5(\Sigma\mathbb{R}P^\infty) \arrow[r] \arrow[d, equal] & H_4(\mathbb{Z}/2) \arrow[d, equal] \\
H_5(\Sigma_3) \arrow[r] & \Gamma_5(\Sigma K(\Sigma_3,1)) \arrow[r] & \pi_5(\Sigma K(\Sigma_3,1)) \arrow[r] & H_4(\Sigma_3)
\end{tikzcd}$$

Here the natural isomorphism $\Gamma_5(\Sigma\mathbb{R}P^\infty) \to \Gamma_5(\Sigma K(\Sigma_3,1))$ follows from the diagram

$$\begin{tikzcd}
L_2\Gamma_2^2(\mathbb{Z}/4 \twoheadrightarrow \mathbb{Z}/2) \arrow[r, equal] \arrow[d] & L_2\Gamma_2^2(\mathbb{Z}/4 \twoheadrightarrow \mathbb{Z}/2) \arrow[d] \\
\Gamma_2^3(\mathbb{Z}/4 \twoheadrightarrow \mathbb{Z}/2,\ \mathbb{Z}/2 \hookrightarrow \mathbb{Z}/4) \arrow[r, equal] \arrow[d] & \Gamma_2^3(\mathbb{Z}/4 \twoheadrightarrow \mathbb{Z}/2,\ \mathbb{Z}/2 \hookrightarrow \mathbb{Z}/12) \arrow[d] \\
\Gamma_5(\Sigma\mathbb{R}P^\infty) \arrow[r] \arrow[d] & \Gamma_5(\Sigma K(\Sigma_3,1)) \arrow[d] \\
L_1\Gamma_2^2(\mathbb{Z}/4 \twoheadrightarrow \mathbb{Z}/2) \arrow[r, equal] & L_1\Gamma_2^2(\mathbb{Z}/4 \twoheadrightarrow \mathbb{Z}/2)
\end{tikzcd}$$

□

5. Relation to K-theory

As we mentioned in the introduction, there is a natural relation between the problem considered and algebraic K-theory. Since the plus-construction $K(G,1) \to K(G,1)^+$ is a homological equivalence, there is a natural weak homotopy equivalence

$$\Sigma K(G,1) \to \Sigma(K(G,1)^+)$$

This defines the natural suspension map:

$$\pi_n(K(G,1)^+) \to \pi_{n+1}(\Sigma(K(G,1)^+)) = \pi_{n+1}(\Sigma K(G,1))$$

for $n \geq 1$.

Given a group G and its maximal perfect normal subgroup $P \lhd G$, one has natural isomorphism $\pi_n(K(P,1)^+) \simeq \pi_n(K(G,1)^+),\ n \geq 2$ since $K(P,1)^+$ is homotopy equivalent to the universal covering space of $K(G,1)^+$.

For a perfect group G, the Whitehead exact sequences form the following commutative diagram:

$$\begin{tikzcd}
H_4(G) \arrow[r] \arrow[d, equal] & \Gamma_2(H_2(G)) \arrow[r] \arrow[d] & \pi_3(K(G,1)^+) \arrow[r, two heads] \arrow[d] & H_3(G) \arrow[d, equal] \\
H_4(G) \arrow[r] & H_2(G) \otimes \mathbb{Z}/2 \arrow[r] & \pi_4(\Sigma K(G,1)) \arrow[r, two heads] & H_3(G)
\end{tikzcd} \tag{5.1}$$

Here we will look at the applications of the following two classical constructions:

1) Let R be a ring and $G = E(R)$, the group of elementary matrices. The group $E(R)$ is perfect and the plus-construction $K(E(R),1)^+$ also denoted $\tilde{K}(R)$, defines the algebraic K-theory of R: $K_n(R) = \pi_n(K(E(R),1)^+),\ n \geq 2$.

2) Let Σ_∞ be the infinite permutation groups and A_∞ is the infinite alternating subgroup. There is the following description of stable homotopy groups of spheres [14]:

$$\pi_n^S = \pi_n(K(\Sigma_\infty,1)^+) = \pi_n(K(A_\infty,1)^+),\ n \geq 2. \tag{5.2}$$

5.1. Let R be a ring. In this case, one has the natural homomorphisms:

$$K_n(R) \to \pi_{n+1}(\Sigma K(E(R),1)),\ n \geq 2.$$

For $n = 2$, clearly one has the natural isomorphism:

$$K_2(R) \simeq H_2(E(R)) \simeq \pi_3(\Sigma K(E(R),1)). \tag{5.3}$$

It is shown in [1] that the map $\Gamma_2(K_2(R)) \to K_3(R)$ factors as

$$\Gamma_2(K_2(R)) \twoheadrightarrow K_2(R) \otimes K_1(\mathbb{Z}) \xrightarrow{\star} K_3(R),$$

where $\star$ is the product in algebraic K-theory: $\star : K_i(S) \otimes K_j(T) \to K_{i+j}(S \otimes T)$. Hence the diagram (5.1) has the following form:

$$\begin{tikzcd}
H_4(E(R)) \arrow[r] \arrow[d, equal] & \Gamma_2(K_2(R)) \arrow[r] \arrow[d] & K_3(R) \arrow[r, two heads] \arrow[d] & H_3(E(R)) \arrow[d, equal] \\
H_4(E(R)) \arrow[r] & K_2(R) \otimes K_1(\mathbb{Z}) \arrow[r] \arrow[ur, "\star"] & \pi_4(\Sigma K(E(R),1)) \arrow[r, two heads] & H_3(E(R))
\end{tikzcd} \tag{5.4}$$

and the natural map

$$K_3(R) \to \pi_4(\Sigma K(E(R),1)) \tag{5.5}$$

is an isomorphism. From equations (5.3) and (5.5) together with the fact that $SL(\mathbb{Z}) = E(\mathbb{Z})$, we have the following:

Theorem 5.1. *The natural homomorphism*

$$K_n(R) \longrightarrow \pi_{n+1}(\Sigma K(E(R),1))$$

is an isomorphism for $n = 2, 3$. *In particular,*

$$\pi_3(\Sigma K(SL(\mathbb{Z}), 1)) \cong K_2(\mathbb{Z}) \cong \mathbb{Z}/2 \text{ and}$$

$$\pi_4(\Sigma K(SL(\mathbb{Z}), 1)) \cong K_3(\mathbb{Z}) \cong \mathbb{Z}/48.$$

□

Remark 5.1. The isomorphism (5.5) and Carlsson construction $F^{E(R)}(S^1)$ gives a way, for an element of $K_3(R)$, to associate an element from $F^{E(R)}(S^1)_3 = E(R) * E(R) * E(R)$ (uniquely modulo $\mathcal{B}F^{E(R)}(S^1)$):

$$\begin{tikzcd}
K_3(R) \arrow[r, rightsquigarrow] \arrow[rr, bend left, "\simeq"] & E(R) * E(R) * E(R) & \frac{\mathcal{Z}F^{E(R)}(S^1)_3}{\mathcal{B}F^{E(R)}(S^1)_3} \arrow[dl, hook] \\
 & \frac{E(R)*E(R)*E(R)}{\mathcal{B}F^{E(R)}(S^1)_3} \arrow[u, "section"] &
\end{tikzcd}.$$

It is interesting to represent in this way known elements from $K_3(R)$ for different rings. For $R = \mathbb{Z}$, $x \in SL(\mathbb{Z}) = E(\mathbb{Z})$, denote by $x^{(1)}, x^{(2)}, x^{(3)}$ the correspondent elements in the free cube $SL(\mathbb{Z}) * SL(\mathbb{Z}) * SL(\mathbb{Z})$. Take the following commuting elements of $SL(\mathbb{Z})$:

$$u = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \qquad v = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

The structure of the element (2.7), diagram (5.4) and well-known facts about structure of $K_2(\mathbb{Z})$ imply that, using the above notations, the element

$$[[u^{(2)}, v^{(3)}], [u^{(1)}, v^{(3)}]]$$

corresponds to the element of order 2 in $K_3(\mathbb{Z})$. It would be interesting to see an element of $SL(\mathbb{Z}) * SL(\mathbb{Z}) * SL(\mathbb{Z})$ which corresponds to the generator of $K_3(\mathbb{Z}) = \mathbb{Z}/48$. □

Consider the case $R = \mathbb{Z}$ and $n = 5$. In this case, $E(\mathbb{Z}) = SL(\mathbb{Z})$ and we have the following commutative diagram with exact horizontal sequences:

$$\begin{tikzcd}
\mathbb{Z} \oplus (\mathbb{Z}/2)^2 \arrow[r, two heads] \arrow[d, equal] & (\mathbb{Z}/2)^3 \arrow[r] \arrow[d, equal] & 0 \arrow[r] \arrow[d, equal] & \mathbb{Z}/2 \arrow[d, equal] & \\
H_5SL(\mathbb{Z}) \arrow[r] \arrow[d, equal] & \Gamma_4(\tilde{K}(\mathbb{Z})) \arrow[r] \arrow[d] & K_4(\mathbb{Z}) \arrow[r] \arrow[d] & \mathbb{Z}/2 \arrow[r, hook] \arrow[d, equal] & \mathbb{Z}/4 \arrow[d, two heads] \\
H_5(SL(\mathbb{Z})) \arrow[r] & \Gamma_5(\Sigma K(SL(\mathbb{Z}), 1)) \arrow[r] & \pi_5(\Sigma K(SL(\mathbb{Z}), 1)) \arrow[r] & \mathbb{Z}/2 \arrow[r, "0"] & \mathbb{Z}/2
\end{tikzcd}$$

and the following commutative diagram:

$$\begin{tikzcd}
(\mathbb{Z}/2)^2 \arrow[r, hook] \arrow[d, equal] & (\mathbb{Z}/2)^3 \arrow[r, two heads] \arrow[d, equal] & \mathbb{Z}/2 \arrow[d, equal] \\
\Gamma_2^2(\Gamma_2(K_2(\mathbb{Z})) \to K_3(\mathbb{Z})) \arrow[r, hook] \arrow[d] & \Gamma_4(\tilde{K}(\mathbb{Z})) \arrow[r, two heads] \arrow[d] & R_2(K_2(\mathbb{Z})) \arrow[d] \\
\pi_4(\Sigma K(SL(\mathbb{Z}),1)) \otimes \mathbb{Z}/2 \arrow[r, hook] \arrow[d, equal] & \Gamma_5(\Sigma K(SL(\mathbb{Z}),1)) \arrow[r] \arrow[d, equal] & Tor(\pi_3(\Sigma K(SL(\mathbb{Z})),1),\mathbb{Z}/2) \arrow[d, equal] \\
\mathbb{Z}/2 \arrow[r, hook] & (\mathbb{Z}/2)^2 \arrow[r, two heads] & \mathbb{Z}/2
\end{tikzcd}$$

Simple analysis shows that the suspension map $\Gamma_4(\tilde{K}(\mathbb{Z})) \to \Gamma_5(\Sigma K(SL(\mathbb{Z}),1))$ is an epimorphism and therefore we have the following theorem:

Theorem 5.2. *The Hurewicz homomorphism*

$$\pi_5(\Sigma K(SL(\mathbb{Z}),1)) \to H_4(SL(\mathbb{Z})) = \mathbb{Z}/2$$

is an isomorphism. □

Remark. Since $K_4(\mathbb{Z}) = 0$, we see that the natural homomorphism

$$K_4(\mathbb{Z}) \to \pi_5(\Sigma K(SL(\mathbb{Z}),1))$$

is not an isomorphism.

5.2. Here we will use (5.2) for certain computations.

Theorem 5.3. *Let A_4 be the 4-th alternating group. Then $\pi_4(\Sigma K(A_4,1)) = \mathbb{Z}/4$.*

Proof. First recall that[3]

$$\begin{array}{llll} H_1(A_4) = \mathbb{Z}/3, & H_2(A_4) = \mathbb{Z}/2, & H_3(A_4) = \mathbb{Z}/6, & H_4(A_4) = 0 \\ H_2(A_\infty) = \mathbb{Z}/6, & H_3(A_\infty) = \mathbb{Z}/12, & \pi_3(\Sigma K(A_4,1)) = \mathbb{Z}/6 & \end{array}$$

Consider the Whitehead exact sequence for the space $\Sigma K(A_4,1)$:

$$\begin{tikzcd}
\Gamma_3(\Sigma K(A_4,1)) \arrow[r, hook] & \pi_4(\Sigma K(A_4,1)) \arrow[r] & H_3(A_4) \arrow[r] \arrow[d, equal] & \Gamma_2(H_1(A_4)) \arrow[r] \arrow[d, equal] & \pi_3(\Sigma K(A_4,1)) \arrow[r, two heads] \arrow[d, equal] & H_2(A_4) \arrow[d, equal] \\
& & \mathbb{Z}/6 \arrow[r] & \mathbb{Z}/3 \arrow[r] & \mathbb{Z}/6 \arrow[r, two heads] & \mathbb{Z}/2
\end{tikzcd}$$

---

[3] These computations were done with the help of HAP-system. The authors thank Graham Ellis for these computations

Since $R_2(\pi_2\Sigma K(A_4,1)) = R_2(\mathbb{Z}/3) = 0$, we have $\Gamma_3(\Sigma K(A_4,1)) = \Gamma_2^2(\mathbb{Z}/3 \hookrightarrow \mathbb{Z}/6)$. It follows from the definition of the functor Γ_2^2 that it is isomorphic to the pushout

$$\begin{tikzcd}
\mathbb{Z}/3\otimes(\mathbb{Z}/3\oplus\mathbb{Z}/2) \arrow[r] \arrow[d] & 0 \arrow[d] \\
\mathbb{Z}/6\otimes(\mathbb{Z}/3\oplus\mathbb{Z}/2) \arrow[r] & \Gamma_2^2(\mathbb{Z}/3\hookrightarrow\mathbb{Z}/6)
\end{tikzcd}$$

That is, $\Gamma_2^2(\mathbb{Z}/3 \hookrightarrow \mathbb{Z}/6) = \mathbb{Z}/2$ and there is the following short exact sequence:

$$0 \to \mathbb{Z}/2 \to \pi_4(\Sigma K(A_4,1)) \to \mathbb{Z}/2 \to 0.$$

We come to the extension problem: is it $\mathbb{Z}/2^2$ or $\mathbb{Z}/4$?

Consider the monomorphism $A_4 \hookrightarrow A_\infty$ and the map between corresponding Whitehead sequences:

$$\begin{tikzcd}
\Gamma_3(\Sigma K(A_4,1)) \arrow[r, hook] \arrow[d] & \pi_4(\Sigma K(A_4,1)) \arrow[r] \arrow[d] & H_3(A_4) \arrow[r] \arrow[d] & \Gamma_2(H_1(A_4)) \arrow[d] \\
\Gamma_3(\Sigma K(A_\infty,1)) \arrow[r] & \pi_4(\Sigma K(A_\infty,1)) \arrow[r] & H_3(A_\infty) \arrow[r] & \Gamma_2(H_1(A_\infty))
\end{tikzcd}$$

which is

$$\begin{tikzcd}
\mathbb{Z}/2 \arrow[r, hook] \arrow[d] & \pi_4(\Sigma K(A_4,1)) \arrow[r] \arrow[d] & \mathbb{Z}/6 \arrow[r] \arrow[d] & \mathbb{Z}/6 \arrow[d] \\
\Gamma_2^2(0\to\mathbb{Z}/6) \arrow[r] & \pi_4(\Sigma K(A_\infty,1)) \arrow[r] & \mathbb{Z}/12 \arrow[r] & 0
\end{tikzcd} \tag{5.6}$$

It is easy to see that $\Gamma_2^2(0 \to \mathbb{Z}/6) = \mathbb{Z}/2$ and that $\Gamma_2^2(\mathbb{Z}/3 \hookrightarrow \mathbb{Z}/6) \to \Gamma_2^2(0 \to \mathbb{Z}/6)$ is an isomorphism. We obtain the following diagram:

$$\begin{tikzcd}
\mathbb{Z}/2 \arrow[r, hook] \arrow[d, equal] & \pi_4(\Sigma K(A_4,1)) \arrow[r, two heads] \arrow[d] & \mathbb{Z}/2 \arrow[d, hook] \\
\mathbb{Z}/2 \arrow[r] & \pi_4(\Sigma K(A_\infty,1)) \arrow[r, two heads] & \mathbb{Z}/12
\end{tikzcd} \tag{5.7}$$

Now we use the isomorphism (5.2). Consider the suspension:

$$K(A_\infty,1)^+ \to \Omega\Sigma K(A_\infty,1)^+ \simeq \Omega\Sigma K(A_\infty,1)$$

and the corresponding map between Whitehead sequences:

$$\begin{tikzcd}
H_4(A_\infty) \arrow[r] \arrow[d, equal] & \Gamma_2(\pi_2^S) \arrow[r] \arrow[d] & \pi_3^S \arrow[r, two heads] \arrow[d] & H_3(A_\infty) \arrow[d, equal] \\
H_4(A_\infty) \arrow[r] & \pi_2^S\otimes\mathbb{Z}/2 \arrow[r] & \pi_4(\Sigma K(A_\infty,1)) \arrow[r] & H_3(A_\infty)
\end{tikzcd}$$

Since

$$\pi_3^S = \mathbb{Z}/24,\ \pi_4^S = 0,$$

we conclude that the Whitehead sequence for $K(A_\infty, 1)^+$ has the following form:

$$\begin{tikzcd}
H_4(A_\infty) \arrow[r, hook] \arrow[d, equal] & \Gamma_2(\pi_2^S) \arrow[r] \arrow[d, equal] & \pi_3^S \arrow[r, two heads] \arrow[d, equal] & H_3(A_\infty) \arrow[d, equal] \\
\mathbb{Z}/2 \arrow[r, hook] & \mathbb{Z}/4 \arrow[r] & \mathbb{Z}/24 \arrow[r, two heads] & \mathbb{Z}/12
\end{tikzcd}$$

We conclude that the map

$$\pi_3^S \to \pi_4(\Sigma K(A_\infty, 1))$$

is an isomorphism and that the map

$$H_4(A_\infty) \to \Gamma_2^2(0 \to \mathbb{Z}/6)$$

is the zero map. The diagram (5.7) has the following form:

$$\begin{tikzcd}
\mathbb{Z}/2 \arrow[r, hook] \arrow[d, equal] & \pi_4(\Sigma K(A_4, 1)) \arrow[r, two heads] \arrow[d] & \mathbb{Z}/2 \arrow[d, hook] \\
\mathbb{Z}/2 \arrow[r, hook] & \mathbb{Z}/24 \arrow[r, two heads] & \mathbb{Z}/12
\end{tikzcd}$$

The result follows. □

REFERENCES

[1] D. Arlettaz: Algebraic K-theory of rings from a topological viewpoint, *Publ. Math.* **44** (2008), 3-84.

[2] H.-J. Baues: *Homotopy type and homology*, Oxford Math. Monographs (1996), Oxford University Press.

[3] H.-J. Baues and D. Conduché: On the tensor algebra of a non-abelian group and application, *K-Theory,* **5** (1992), 531–554.

[4] H.-J. Baues and P. Goerss: A homotopy operation spectral sequence for the computation of homotopy groups, *Topology* **39** (2000), 161-192.

[5] L. Breen: On the functorial homology of abelian groups, *J. Pure Appl. Alg.* **142** (1999), 199-237.

[6] R. Brown and J.-L. Loday: Van Kampen theorems for diagrams of spaces, *Topology* **26** (1987), 311–335.

[7] G. Carlsson: A simplicial group construction for balanced products, *Topology* **23** (1985), 85–89.

[8] F. R. Cohen, J. C. Moore, and J. A. Neisendorfer: Torsion in homotopy groups, *Ann. of Math.* (2) **109** (1979), 121–168.

[9] A. Dold and D. Puppe: Homologie nicht-additiver Funtoren; Anwendugen. *Ann. Inst. Fourier* **11** (1961)

[10] S. Eilenberg and S. Mac Lane: On the groups $H(\pi, n)$, II: Methods of computation, *Ann. Math.* **60**, (1954), 49–139.

[11] G. Ellis and R. Mikhailov: A colimit of classifying spaces, preprint http://arxiv.org/abs/0804.3581

[12] J. C. Harris and N. J. Kuhn: Stable decompositions of classifying spaces of finite abelian p-groups. *Math. Proc. Cambridge Philos. Soc.* **103** (1988) 427–449.

[13] J. A. Neisendorfer: Primary homotopy theory. *Mem. Amer. Math. Soc.* 25 (1980), no. 232, iv+67 pp.

[14] S. Priddy: On $\Omega^\infty S^\infty$ and the infinite symmetric group, *Algebraic Topology, Proc. Sympos. Pure Math.* **22**, 217-220 (1971).

[15] D. G. Quillen: Spectral sequences of a double semi-simplicial group. *Topology* **5** (1966) 155–157.
[16] H. Toda: Composition methods in homotopy groups of spheres, Princeton Univ. Press, 1962.
[17] H. Toda: Order of the identity class of a suspension space. *Ann. Math.* **78** (1963) 300–325.
[18] J. H. C. Whitehead: The homotopy type of special kind of polyhedron, *Annals de la Soc. Polon. de Math.* **21** (1948), 51–110.
[19] J. H. C. Whitehead: A certain exact sequence, *Ann. Math.* **52** (1950), 51–110.
[20] G. W. Whitehead: On spaces with vanishing low-dimensional homotopy groups, *proc. Nat. Acd. Sci.* **34** (1948), 207–211.
[21] J. Wu: Combinatorial description of homotopy groups of certain spaces, *Math. Proc. Camb. Phyl. Soc.* 130, (2001), 489–513.
[22] J. Wu: Homotopy theory of the suspensions of the projective plane *Memoirs AMS* 162 (2003), No. 769.

Steklov Mathematical Institute, Gubkina 8, Moscow, Russia 119991
E-mail address: `romanvm@mi.ras.ru`

Department of Mathematics, National University of Singapore, 2 Science Drive 2 Singapore 117542
E-mail address: `matwuj@nus.edu.sg`
URL: `www.math.nus.edu.sg/~matwujie`